

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
22 February 2001 (22.02.2001)

PCT

(10) International Publication Number  
WO 01/12668 A1

(51) International Patent Classification<sup>7</sup>: C07K 14/47,  
7/08, A61K 38/17, A61P 31/04

(74) Agent: HASH, Stephen, M.; Fulbright & Jaworski,  
L.L.P., Suite 2400, 600 Congress Avenue, Austin, TX  
78701 (US).

(21) International Application Number: PCT/US00/22781

(22) International Filing Date: 18 August 2000 (18.08.2000)

(25) Filing Language: English

(26) Publication Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,  
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,  
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,  
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO,  
NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR,  
TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(30) Priority Data:  
60/149,886 18 August 1999 (18.08.1999) US

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,  
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicants: UNIVERSITY OF IOWA RESEARCH  
FOUNDATION [US/US]; 100 Oakdale Campus  
#214-TIC, Iowa City, IA 52242-5000 (US). THE RE-  
GENTS OF THE UNIVERSITY OF CALIFORNIA  
[US/US]; 12th Floor, 1111 Franklin Street, Oakland, CA  
94607 (US).

**Published:**

- With international search report.
- Before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments.

(72) Inventors: TACK, Brian, E.; 213 McLean Street, Iowa  
City, IA 55246 (US). MCCRAY, Paul; 534 Park Street,  
Iowa City, IA 52240 (US). WELSH, Michael; 3460 564th  
Street, Riverside, IA 52327 (US). TRAVIS, Sue, M.; 405  
S. Summit, Iowa City, IA 52240 (US). LEHRER, Robert;  
..

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.



WO 01/12668 A1

(54) Title: CATHELICIDIN-DERIVED PEPTIDES WITH BROAD SPECTRUM ANTIMICROBIAL ACTIVITY

(57) Abstract: The present invention relates to the use of antimicrobial peptides in the inhibition of microbial growth and proliferation. Novel antimicrobial truncated peptides are disclosed which are based upon SMAP 29 and RCAP 18, but which contain a lesser number of amino acid residues yet still retain bactericidal activity. In addition, synthetic peptides based upon the SMAP 29 protein are disclosed which have fewer amino acid residues and include substitutions yet retain substantial activity. The invention also relates to a method of inhibiting microbial growth by administering an effective amount of a peptide in accordance with the invention, or by combining the peptides with other antimicrobial agents or antibiotics.

BEST AVAILABLE COPY

## CATHELICIDIN-DERIVED PEPTIDES WITH BROAD SPECTRUM ANTIMICROBIAL ACTIVITY

**BACKGROUND OF THE INVENTION**

The government may have rights in the present invention pursuant to grant number NIH P50 HL-61234-01 from the National Institutes of Health, Cystic Fibrosis Foundation

5

**1. Field of the Invention**

The present invention relates generally to the fields of molecular biology and microbiology. More particularly, it concerns the use of antimicrobial peptides for the inhibition of microbial growth.

10

**2. Description of Related Art**

The first antibiotics were used clinically in the 1940s and 1950s, and their use has been increasing significantly since this period. Although an invaluable advance, antibiotic and antimicrobial therapy suffers from several problems, particularly when strains of various bacteria appear that are resistant to antibiotics. Interestingly, bacteria resistant to streptomycin were isolated about a year after this antibiotic was introduced (Waksman, 1945).

The development of antibiotic resistance is a serious and life-threatening event of worldwide importance. For example, strains of *Staphylococcus* are known that are immune to all antibiotics except one (Travis, 1994). Such bacteria often cause fatal hospital infections. Among other drug resistant organisms are: pneumococci that cause pneumonia and meningitis; *Cryptosporidium* and *E. coli* that cause diarrhea; and enterococci that cause blood-stream, surgical wound and urinary tract infections (Berkelman *et al.*, 1994). The danger is further compounded by the fact that antibiotic and antimicrobial resistance may be spread vertically and horizontally by plasmids and transposons.

Davies (1986) described seven basic biochemical mechanisms for naturally-occurring antibiotic resistance: (1) alteration (inactivation) of the antibiotic; (2) alteration

of the target site; (3) blockage in the transport of the antibiotic; (4) by-pass of the antibiotic sensitive-step (replacement); (5) increase in the level of the inhibited enzyme (titration of drug); (6) sparing the antibiotic-sensitive step by endogenous or exogenous product; and (7) production of a metabolite that antagonizes action of inhibitor.

5

Antimicrobial peptides have been isolated from plants, insects, fish, amphibia, birds, and mammals (Gallo, 1998; Ganz & Lehrer, 1998). Although previously considered an evolutionarily ancient system of immune protection with little relevance beyond minimal primary protection, recent developments have found that mammalian  
10 cells express these peptide antibiotics during inflammatory events such as wound repair, contact dermatitis and psoriasis (From Nilsson, 1999). These peptides are apparently a primary component of innate host protection against microbial pathogenesis functioning to create pores in the cytoplasmic membrane of microorganisms (Oren *et al.*, 1998). Furthermore, antimicrobial peptides also act on animal cells by stimulating them to  
15 change behaviors such as syndecan expression, chemotaxis, and chloride secretion (Gallo, 1998). After contact with microorganisms, vertebrate skin, trachea and tongue epithelia are rich sources of peptide antibiotics, which may explain the unexpected resistance of these tissues to infection (Russell *et al.* 1996).

20 The present invention seeks to overcome these and other drawbacks inherent in the prior art by providing new compositions, combined compositions, methods and kits, for use in reducing resistance to antimicrobials and antibiotics and for treating infections.

### SUMMARY OF THE INVENTION

25

The present invention provides new methods, combined compositions and kits, for use in inhibiting microbial growth and proliferation, reducing resistance to antimicrobials, and providing novel antibiotics for treating infections. The invention rests in the surprising use of one or more antimicrobial peptides alone, or in conjunction  
30 with an antimicrobial agent or antibiotic in the control of microbial growth and or proliferation.

The invention therefore encompasses methods, compositions, and kits that relate to an antimicrobial peptide. One embodiment thus represents an isolated antimicrobial peptide comprising the amino acid sequence: KNLRRIIRKIIHIKKYG-NH<sub>2</sub> (SEQ ID NO: 1), KNIRRIIRKIIHIKKYG-NH<sub>2</sub> (SEQ ID NO: 6), KNIRRIIRKIIHIKKYG (SEQ ID NO: 7), KNLRRIIRKIIHIKKYG (SEQ ID NO: 8), NLRRIIRKIIHIKKY (SEQ ID NO 9), NIRRIIRKIIHIKKY (SEQ ID NO: 10), LRRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 11), LRRIIRKIIHIKK (SEQ ID NO: 12), IRRRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 13), IRRRIIRKIIHIKK (SEQ ID NO: 14), LRRIIRKIIHIK-NH<sub>2</sub> (SEQ ID NO: 15), RRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 16), RRIIRKIIHIK-NH<sub>2</sub> (SEQ ID NO: 17), GLRKRLRKFRNKIKEKLKKIG (SEQ ID NO: 19), KRLRKFRNKIKEKLKKIG (SEQ ID NO: 20), RKRLRKFRNKIKEKLKKIGQKI (SEQ ID NO: 21), LRKFRNKIKEKLKKIGQKI (SEQ ID NO: 22), LRKFRNKIKEKLKKIGQKIQG (SEQ ID NO: 23), RKFRNKIKEKLKKIG (SEQ ID NO: 24), KIKEKLKKIGQKIQG (SEQ ID NO: 25), KIKEKLKKIGQKIQGLL (SEQ ID NO: 26), RGLRRLGRKIAHGVKKYGPTVLRIRIA-NH<sub>2</sub> (SEQ ID NO. 27), or KNLRRIIRKIIHIKKYGPTILRIIRIIG-NH<sub>2</sub> (SEQ ID NO. 28).

An additional embodiment would consist of a pharmaceutical composition wherein said composition comprises the antimicrobial peptide comprising the amino acid sequence: KNLRRIIRKIIHIKKYG-NH<sub>2</sub> (SEQ ID NO: 1) , KNIRRIIRKIIHIKKYG-NH<sub>2</sub> (SEQ ID NO:6), KNIRRIIRKIIHIKKYG (SEQ ID NO: 7), KNLRRIIRKIIHIKKYG (SEQ ID NO: 8), NLRRIIRKIIHIKKY (SEQ ID NO 9), NIRRIIRKIIHIKKY (SEQ ID NO: 10), LRRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 11), LRRIIRKIIHIKK (SEQ ID NO: 12), IRRRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 13), IRRRIIRKIIHIKK (SEQ ID NO: 14), LRRIIRKIIHIK-NH<sub>2</sub> (SEQ ID NO: 15), RRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 16), RRIIRKIIHIK-NH<sub>2</sub> (SEQ ID NO: 17), GLRKRLRKFRNKIKEKLKKIG (SEQ ID NO: 19), KRLRKFRNKIKEKLKKIG (SEQ ID NO: 20), RKRLRKFRNKIKEKLKKIGQKI (SEQ ID NO: 21), LRKFRNKIKEKLKKIGQKI (SEQ ID NO: 22), LRKFRNKIKEKLKKIGQKIQG (SEQ ID NO: 23), RKFRNKIKEKLKKIG (SEQ ID NO: 24), KIKEKLKKIGQKIQG



(SEQ ID NO: 25), KIKEKLKKIGQKIQGLL (SEQ ID NO: 26)  
 RGLRRLGRKIAHGVKKYGPTVLRIRIA-NH<sub>2</sub> (SEQ ID NO. 27), or  
 KNLRRIIRKIIHIKKYGPTILRIIRIG-NH<sub>2</sub> (SEQ ID NO. 28), and a pharmaceutically  
 acceptable carrier.

5

In a further embodiment of the invention, an antimicrobial peptide will be  
 introduced into an environment, including but not limited to a host, in order to inhibit the  
 growth and/or proliferation of microbial organisms. Such an introduction envisions that  
 the microbial organism will be contacted by the antimicrobial peptide, and as a result of  
 10 this contact, the growth and or proliferation of the microbial organism will be inhibited.  
 A preferred embodiment is a method of inhibiting microbial growth in an environment  
 capable of sustaining such growth comprising administering to said environment the  
 antimicrobial peptide comprising the amino acid sequence: KNLRRIIRKIIHIKKYG-  
 NH<sub>2</sub> (SEQ ID NO: 1), KNIRRIIRKIIHIKKYG-NH<sub>2</sub> (SEQ ID NO: 6),  
 15 KNIRRIIRKIIHIKKYG (SEQ ID NO: 7), KNLRRIIRKIIHIKKYG (SEQ ID NO: 8),  
 NLRIIRKIIHIKKY (SEQ ID NO 9), NIRRIIRKIIHIKKY (SEQ ID NO: 10),  
 LRRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 11), LRRIIRKIIHIKK (SEQ ID NO: 12),  
 IRRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 13), IRRIIRKIIHIKK (SEQ ID NO: 14),  
 LRRIIRKIIHIK-NH<sub>2</sub> (SEQ ID NO: 15), RRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 16),  
 20 RRIIRKIIHIK-NH<sub>2</sub> (SEQ ID NO: 17),  
 GLRKRLRKFRNKIKEKLKKIGQKIQGLLPKLAPRTDY (SEQ ID NO: 18),  
 GLRKRLRKFRNKIKEKLKKIG (SEQ ID NO: 19), KRLRKFRNKIKEKLKKIG (SEQ  
 ID NO: 20), RKRLRKFRNKIKEKLKKIGQKI (SEQ ID NO: 21),  
 LRKFRNKIKEKLKKIGQKI (SEQ ID NO: 22), LRKFRNKIKEKLKKIGQKIQG  
 25 (SEQ ID NO: 23), RKFRNKIKEKLKKIG (SEQ ID NO: 24), KIKEKLKKIGQKIQG  
 (SEQ ID NO: 25), KIKEKLKKIGQKIQGLL (SEQ ID NO: 26)  
 RGLRRLGRKIAHGVKKYGPTVLRIRIA-NH<sub>2</sub> (SEQ ID NO. 27), or  
 KNLRRIIRKIIHIKKYGPTILRIIRIG-NH<sub>2</sub> (SEQ ID NO. 28)..

30

Such a method may further consist of administering an antimicrobial peptide in a  
 pharmaceutically acceptable carrier and/or in combination with a second antimicrobial

agent or antibiotic. Such antimicrobial agents or antibiotics may include but are not limited to the peptides listed in SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20, SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25 SEQ ID NO: 26, SEQ ID NO: 27 or SEQ ID NO: 28 or a protein synthesis inhibitor, a cell wall growth inhibitor, a cell membrane synthesis inhibitor, a nucleic acid synthesis inhibitor, and a competitive inhibitor.

An additional embodiment would consist of a method of inhibiting microbial growth in a host, comprising administering to said host the antimicrobial peptide comprising the amino acid sequence: KNLRRIIRKIIHIKKYG-NH<sub>2</sub> (SEQ ID NO: 1), RGLRRLGRKIAHGVKKYGPTVLRIRIAG (SEQ ID NO: 2), KIAHGVKKYGPTVLRIRIAG (SEQ ID NO: 3), LGRKIAHGVKKYGPTVLRIRI (SEQ ID NO: 4), RGLRRLGRKIAHGVKKYG (SEQ ID NO: 5), KNIRRIIRKIIHIKKYG-NH<sub>2</sub> (SEQ ID NO: 6), KNIRRIIRKIIHIKKYG (SEQ ID NO: 7), KNLRRIIRKIIHIKKYG (SEQ ID NO: 8), NLRRIIRKIIHIKKY (SEQ ID NO: 9), NIRRIIRKIIHIKKY (SEQ ID NO: 10), LRRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 11), LRRIIRKIIHIKK (SEQ ID NO: 12), IIRRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 13), IIRRIIRKIIHIKK (SEQ ID NO: 14), LRRIIRKIIHIK-NH<sub>2</sub> (SEQ ID NO: 15), RRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 16), RRIIRKIIHIK-NH<sub>2</sub> (SEQ ID NO: 17), GLRKRLRKFRNKIKEKLKKIGQKIQGLLPKLAPRTDY (SEQ ID NO: 18), GLRKRLRKFRNKIKEKLKKIG (SEQ ID NO: 19), KRLRKFRNKIKEKLKKIG (SEQ ID NO: 20), RKRLRKFRNKIKEKLKKIGQKI (SEQ ID NO: 21), LRKFRNKIKEKLKKIGQKI (SEQ ID NO: 22), LRKFRNKIKEKLKKIGQKIQG (SEQ ID NO: 23), RKFRNKIKEKLKKIG (SEQ ID NO: 24), KIKEKLKKIGQKIQG (SEQ ID NO: 25), KIKEKLKKIGQKIQGLL (SEQ ID NO: 26) RGLRRLGRKIAHGVKKYGPTVLRIRIA-NH<sub>2</sub> (SEQ ID NO. 27), or KNLRRIIRKIIHIKKYGPTILRIIRIIG-NH<sub>2</sub> (SEQ ID NO. 28).

The microorganism, *e.g.*, bacterium, yeast, or population thereof, may be contacted either *in vitro* or *in vivo*. Contacting *in vivo* may be achieved by administering

to an animal (including a human patient) that has, or is suspected to have a microbial or bacterial infection, a therapeutically effective amount of pharmacologically acceptable antimicrobial peptide formulation alone or in combination with a therapeutic amount of a pharmacologically acceptable formulation of a second agent. The invention may thus be employed to treat both systemic and localized microbial, bacterial, and fungal infections by introducing the agent or agents into the general circulation or by applying the combination, *e.g.*, topically to a specific site, such as a wound or burn, or to the eye, ear or other site of infection.

10       An “effective amount of an antimicrobial agent or peptide” means an amount, or dose, within the range required to inhibit bacterial growth and/or proliferation. Such ranges would be readily determinable to those of skill in the art depending upon the use to which the peptide is to be applied. An “effective amount of an antimicrobial agent or antibiotic” means an amount, or dose, within the range normally given or prescribed.

15       Such ranges are well established in routine clinical practice and will thus be readily determinable to those of skill in the art. Doses may be measured by total amount given or by concentration. Ten, 20, 50, 100, 500 and 1000 µg/ml solutions all are appropriate for treatment. Total doses of 100, 500, 1000, 2000, 5000 and 10,000 µg also appear to be appropriate doses. Appropriate oral and parenteral doses and treatment regimens are

20       further detailed herein in Table 5 and Table 6.

As this invention provides for enhanced microbial and/or bacterial killing, it will be appreciated that effective amounts of an antimicrobial agent or antibiotic may be used that are lower than the standard doses previously recommended when the antimicrobial or antibiotic is combined with an antimicrobial peptide. It is further envisioned that the antimicrobial peptide may be used in combination with other antimicrobial agents or antibiotics for a variety of purposes. These purposes include but are not limited to: enhancing the activity of an agent or antibiotic, allowing for a lower dose of an agent or antibiotic due to toxicity or dosing concerns relating to the second agent or antibiotic, enhancing the activity of agents or antibiotics against strains that have previously exhibited resistance to an agent or antibiotic or providing an additional agent in

25       

30

individuals whose immune system is damaged or compromised and are thus unable to mount an effective immune response.

Where a combination of an antimicrobial peptide and one or more antimicrobial agents or antibiotics is contemplated, it is envisioned that the antimicrobial peptide and the second agent or antibiotic may be delivered either simultaneously or either of the agents may be administered prior to the administration of the other. It is envisioned that staggered administration might weaken the microbial organism and increase the efficacy of the additional agent.

In a preferred embodiment of the invention, an antimicrobial peptide will be used alone or in combination with one or more additional antimicrobial agents or antibiotics in the treatment of microbial strains previously determined to be resistant to one or more methods of treatment. It is envisioned that this method will comprise inhibiting the growth of drug-resistant microbial strains comprising administering to an environment capable of sustaining such growth an antimicrobial peptide comprising the amino acid sequence:

KNLRRRIIRKIIHIIKKYG-NH<sub>2</sub> (SEQ ID NO: 1),  
 RGLRRLGRKIAHGVKKYGPTVLRIIRIAG (SEQ ID NO: 2),  
 KIAHGVKKYGPTVLRIIRIAG (SEQ ID NO: 3), LGRKIAHGVKKYGPTVLRII (SEQ ID NO: 4),  
 RGLRRLGRKIAHGVKKYG (SEQ ID NO: 5), KNIRRIIRKIIHIIKKYG-NH<sub>2</sub> (SEQ ID NO: 6),  
 KNIRRIIRKIIHIIKKYG (SEQ ID NO: 7), KNLRRRIIRKIIHIIKKYG (SEQ ID NO: 8),  
 NLRRIIRKIIHIIKKY (SEQ ID NO: 9), NIRRIIRKIIHIIKKY (SEQ ID NO: 10),  
 LRRRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 11), LRRRIIRKIIHIIKK (SEQ ID NO: 12),  
 IRRRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 13), IRRRIIRKIIHIIKK (SEQ ID NO: 14),  
 LRRRIIRKIIHIIK-NH<sub>2</sub> (SEQ ID NO: 15), RRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 16),  
 RRIIRKIIHIIK-NH<sub>2</sub> (SEQ ID NO: 17), GLRKRLRKFRNKIKEKLKKIGQKIQGLLPKLAPRTDY (SEQ ID NO: 18),  
 GLRKRLRKFRNKIKEKLKKIG (SEQ ID NO: 19), KRLRKFRNKIKEKLKKIG (SEQ ID NO: 20),  
 RKRLRKFRNKIKEKLKKIGQKI (SEQ ID NO: 21), LRKFRNKIKEKLKKIGQKI (SEQ ID NO: 22),  
 LRKFRNKIKEKLKKIGQKI (SEQ ID NO: 23), RKFRNKIKEKLKKIG (SEQ ID NO: 24), KIKEKLKKIGQKI (SEQ ID NO: 25)

NO: 25), KIKEKLKKIGQKIQGLL (SEQ ID NO: 26)  
 RGLRRLGRKIAHGVKKYGPTVLRIRIA-NH<sub>2</sub> (SEQ ID NO: 27), or  
 KNLRRIIRKIIHIKKYGPTILRIIRIG-NH<sub>2</sub> (SEQ ID NO: 28)..

5 Pharmaceutically acceptable compositions may be formulated such that resistant strains may be treated in a host either *ex vivo* or *in vivo* depending upon the requisite circumstances. A list of resistant strains against which an antimicrobial peptide might be employed includes, but is not limited to: *Pseudomonas aeruginosa*, *Burkholderia cepacia*, *Alcaligenes*, *Xanthamonas*, *Listeria monocytogenes*, *Staphylococcus aureus* and  
 10 *Escherichia coli*

In a preferred embodiment of the present invention, the antimicrobial peptides are useful alone or in combination with other antimicrobial agents or antibiotics in the treatment of cystic fibrosis or other diseases of the respiratory system. In the setting of  
 15 cystic fibrosis, the peptides may be used either prophylactically or therapeutically. Because a variety of bacterial strains cause infections associated with cystic fibrosis the broad spectrum properties of antimicrobial peptides make them well suited for use in the prevention of infection associated with the disease.

20 A further embodiment of the invention envisions a nucleic acid molecule encoding the antimicrobial peptide comprising the amino acid sequence: KNLRRIIRKIIHIKKYG-NH<sub>2</sub> (SEQ ID NO: 1), KNIRRIIRKIIHIKKYG-NH<sub>2</sub> (SEQ ID NO: 6), KNIRRIIRKIIHIKKYG (SEQ ID NO: 7), KNLRRIIRKIIHIKKYG (SEQ ID NO: 8), NLRRIIRKIIHIKKY (SEQ ID NO 9), NIRRIIRKIIHIKKY (SEQ ID NO: 10),  
 25 LRRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 11), LRRIIRKIIHIKK (SEQ ID NO: 12), IRRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 13), IRRIIRKIIHIKK (SEQ ID NO: 14), LRRIIRKIIHIK-NH<sub>2</sub> (SEQ ID NO: 15), RRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 16), RRIIRKIIHIK-NH<sub>2</sub> (SEQ ID NO: 17), GLRKRLRKFRNKIKEKLKKIG (SEQ ID NO: 19), KRLRKFRNKIKEKLKKIG (SEQ ID NO: 20), RKRLRKFRNKIKEKLKKIGQKI  
 30 (SEQ ID NO: 21), LRKFRNKIKEKLKKIGQKI (SEQ ID NO: 22), LRKFRNKIKEKLKKIGQKIQG (SEQ ID NO: 23), RKFRNKIKEKLKKIG (SEQ ID



NO: 24), KIKEKLKKIGQKIQG (SEQ ID NO: 25), KIKEKLKKIGQKIQGLL (SEQ ID NO: 26) RGLRRLGRKIAHGVKKYGPTVLRIIRIA-NH<sub>2</sub> (SEQ ID NO. 27), or KNLRRIIRKIIHIKKYGPTILRIIRIIG-NH<sub>2</sub> (SEQ ID NO. 28)..

5 It is envisioned that uses of this nucleic acid sequence could include, but are not limited to, creation of degenerate probes for the detection of further antimicrobial peptide species, use in gene therapy or in the creation of fusion constructs linking the antimicrobial peptides of the instant invention to other proteins.

10 A further embodiment consists of a kit for use in inhibiting microbial growth in a host comprising an antimicrobial peptide comprising the amino acid sequence: KNLRRIIRKIIHIKKYG-NH<sub>2</sub>, (SEQ ID NO: 1), RGLRRLGRKIAHGVKKYGPTVLRIIRIAG (SEQ ID NO: 2), KIAHGVKKYGPTVLRIIRIAG (SEQ ID NO 3), LGRKIAHGVKKYGPTVLRII (SEQ ID NO: 4), RGLRRLGRKIAHGVKKYG (SEQ ID NO: 5), KNIRRIIRKIIHIKKYG-NH<sub>2</sub>, (SEQ ID NO: 6), KNIRRIIRKIIHIKKYG (SEQ ID NO: 7), KNLRRIIRKIIHIKKYG (SEQ ID NO: 8), NLRRIIRKIIHIKKY (SEQ ID NO 9), NIRRIIRKIIHIKKY (SEQ ID NO: 10), LRRIIRKIIHIKK-NH<sub>2</sub>, (SEQ ID NO: 11), LRRIIRKIIHIKK (SEQ ID NO: 12), IRRIRKIIHIKK-NH<sub>2</sub>, (SEQ ID NO: 13), IRRIRKIIHIKK (SEQ ID NO: 14), LRRIIRKIIHIK-NH<sub>2</sub>, (SEQ ID NO: 15), RRIIRKIIHIKK-NH<sub>2</sub>, (SEQ ID NO: 16), RRIIRKIIHIK-NH<sub>2</sub>, (SEQ ID NO: 17), GLRKRLRKFRNKIKEKLKKIGQKIQGLLPKLAPRTDY (SEQ ID NO: 18), GLRKRLRKFRNKIKEKLKKIG (SEQ ID NO: 19), KRLRKFRNKIKEKLKKIG (SEQ ID NO: 20), RKRLRKFRNKIKEKLKKIGQKI (SEQ ID NO: 21), LRKFRNKIKEKLKKIGQKI (SEQ ID NO: 22) LRKFRNKIKEKLKKIGQKIQG (SEQ ID NO: 23), RKFRNKIKEKLKKIG (SEQ ID NO: 24), KIKEKLKKIGQKIQG (SEQ ID NO: 25), KIKEKLKKIGQKIQGLL (SEQ ID NO: 26) RGLRRLGRKIAHGVKKYGPTVLRIIRIA-NH<sub>2</sub> (SEQ ID NO. 27), or KNLRRIIRKIIHIKKYGPTILRIIRIIG-NH<sub>2</sub> (SEQ ID NO. 28). in a suitable container.

25

30 In an additional embodiment, a kit may contain the antimicrobial peptide and a second antimicrobial agent. The second antimicrobial agent may be selected from the group

consisting of a protein synthesis inhibitor, a cell wall growth inhibitor, a cell membrane synthesis inhibitor, a nucleic acid synthesis inhibitor, and a competitive inhibitor.

For the purpose of the instant invention, the term microbe or microbial is considered to encompass, but not be limited to a microscopic pathogen, such as, for example, a virus, virion, bacteria, rickettsial, helminth, nematode, fungi, protozoan or other invasive pathogen.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

FIG. 1: Antimicrobial activity measured with luminescence assay. Relationship between viability (cfu, o) and luminescence (●) of *P. aeruginosa* PAO1 containing pBBR1MCS-5. Bacteria were incubated with the indicated concentration of LL37 for 5 hr, then luminescence was measured and surviving organisms were plated and counted. Studies were performed at an ionic strength of 25 mM (A) and 175 mM (B). Values are percent of control in absence of LL37. Symbols indicate mean  $\pm$  sem, n = 4.

FIGS. 2A and 2B: Effect of ionic strength of antimicrobial activity of CAP18 (FIG. 2A) and SMAP peptides (FIG. 2B). *P. aeruginosa* was incubated with the indicated concentrations of antimicrobial peptide in the standard assay buffer at 25 mM ionic strength (o), or with 150 mM NaCl added (●). Luminescence was measured after 5 hr. Values are percent of control in the absence of antimicrobial peptide; controls were determined for each salt concentration. Symbols indicate mean  $\pm$  range, n = 2; in most cases, the error bars are covered by the symbol.

FIGS. 3A and 3B: CD spectra of CAP 18 (FIG. 3A) and SMAP peptides (FIG. 3B). Spectra of selected peptides were recorded in 0.05 M phosphate buffer pH 7.0 (solid

curve), 0.1% LPS (dotted curve) and in a 40% solution of trifluoroethanol (dashed curve) at peptide concentrations of 0.2 mg/ml.

FIGS. 4A and 4B: Hydrophobicity gradients of cathelicidin peptides. FIG. 4A. The hydrophobicity of each 11-residue window was calculated as described by Eisenberg (1986), and a line was fit by linear regression. The more hydrophobic regions have more negative hydrophobicity values. FIG. 4B. Relationship between antipseudomonal potency (PAOI) and the slope of the hydrophobicity gradient across the peptide.

FIGS. 5A and 5B: Effects of CAP 18 (FIG. 5A) and SMAP 29 (FIG. 5B) peptides on human red blood cells. Erythrocytes were incubated with the indicated concentration of each peptide and hemolysis was measured as discussed in the Examples. Values are percent of control in the absence of antimicrobial peptide. Bars indicate mean  $\pm$  range, n = 2; in some cases, the error bars are covered by the symbols.

15

FIG. 6: Postmortem examination of the lungs of lambs infected with an ovine pneumonic isolate of *Pasturella haemolytica* and subsequently treated with SMAP 29. In infected lambs that did not receive SMAP 29 (Group 3), the pathologic gross lesions were higher than in the SMAP 29 treated group (Group 4) and the scores for microscopic lesions were also higher for the untreated group. SMAP 29 alone was well tolerated and did not induce significant changes at the pulmonary deposition site (Group 2).

20

FIG. 7: Major Conformer A of Smap29 by 2D NMR.

FIG. 8: Major Conformer B of Smap29 by 2D NMR.

FIG. 9: Antimicrobial activity of SMAP-28 in 100mM NaCl (Gram-positive bacteria).

25

FIG. 10: Antimicrobial activity of SMAP-28 in 100mM NaCl (Gram-negative bacteria)

### DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

Although molecules with antibiotic properties have revolutionized medicine, increasingly their unrestrained use has led to the development of resistance in pathogenic strains. Therefore, in order to maintain the present standards of public health, new methods of controlling microbial infection must be devised.

Antimicrobial peptides of higher eukaryotes, though long recognized as components of the innate immune system, were initially considered primitive and of little clinical significance. However, the relative simplicity of these peptides belies their importance, not only in the prevention of primary microbial infection, but also in subsequent immunomodulation (Bowman, 1995). Further, the small size of the molecules suggests a decreased sensitivity to many of the mechanisms of microbial resistance.

Antimicrobial peptides are generally lethal to bacteria and some fungi. They exhibit differential toxicity towards mammalian cells (Hwang *et al.*, 1998). While the mechanism of this action is not definitively known, it is nonetheless believed that the peptides interact with the lipid bilayer and may thus compromise the integrity of the bacterial membrane (Hwang P. et al. 1998).

SMAP 29 is an ovine antimicrobial peptide of the cathelicidin family originally identified through 3' RACE analysis of sheep bone marrow RNA (Mahoney, 1995; Bagella, 1995). RCAP 18 is a lupine antimicrobial peptide of the cathelicidin family originally identified from granulocytes (Hirata, 1994). The present invention arises, in one aspect, from the discovery of methods and compositions which capitalize upon unanticipated and unrecognized characteristics of these molecules.

It has been determined that the SMAP 29 and RCAP 18 peptides are surprisingly effective against bacterial strains previously determined to be resistant to one or more methods of control. More particularly, SMAP 29 compositions are capable of controlling infections by *Pseudomonas aeruginosa*, *Alcaligenes xylosoxidans*, and

*Stenotrophomonas maltophilia*, all strains previously exhibiting resistance to one or more conventional antibiotics. SMAP 29 and RCAP 18 are envisioned to be useful either alone or in combination with other antimicrobial agents or antibiotics in the control of, not only microbial strains normally susceptible to traditional antibiotics, but also those  
5 which have a recognized resistance to conventional therapies.

Further embodiments of the invention relate to truncated versions of the SMAP 29 and RCAP 18 peptides, which retain the bacteriostatic properties. It has been determined that the removal of certain terminal residues of these antimicrobial peptides does not  
10 profoundly effect the antimicrobial properties of the peptide. Thus, truncated peptides have been synthesized to determine what regions of the molecule are critical to antimicrobial function. Truncations include, for example, removal of the 8 C-terminal residues of the SMAP29 molecule (SEQ ID NO: 3), removal of the 4 C-terminal and the  
15 N-terminal residues of the SMAP 29 molecule (SEQ ID NO: 4), and removal of the 11 N-terminal residues of the SMAP 29 molecule (SEQ ID NO: 5). Truncated RCAP 18 peptides with 18-22 amino acids retained activity against Gram-negative bacteria in high salt, but generally lost considerable activity against MRSA under these conditions.

While truncated peptides have utility in determining the active regions of the full  
20 peptide, they also are appropriate themselves for use in the control of microbial growth and proliferation. It is envisioned that those uses determined to be appropriate for the full SMAP29 and RCAP 18 molecules also are appropriate for the truncated versions of the molecules synthesized in specific embodiments of the invention.

25 As the important residues of the molecule were elicited, substitutions were envisioned which would maintain the bacteriostatic properties of such a peptide. Thus, this invention embodies synthesized peptides with a configuration related to that of SMAP 29 or RCAP 18, but possessing alternate structural properties deemed to maintain its antimicrobial characteristics. Alternate constructs, as set forth in SEQ ID NO: 1, 6-17  
30 and 19-28, possess antimicrobial properties similar, but analogous or potentially superior



to those exhibited by the SMAP 29 peptide or RCAP 18 peptide. There also is evidence that SMAPs and RCAPs can synergize with complement to increase cell killing.

The antibacterial potency of the full-length antimicrobial peptides correlated  
5 directly with the hydrophobicity gradient along their backbone, inversely with their relative abundance of anionic residues, and not with the extent of helix formation in trifluoroethanol. The impressive antipseudomonal and antistaphylococcal properties of SMAP 29 and RCAP 18, and the retention of activity in high salt solutions, suggests they, or derivatives thereof, will be effective therapeutic agents for bacterial infections on  
10 airway surfaces, for example, in patients with cystic fibrosis.

The antimicrobial potency of the disclosed molecules also translates to the ability to significantly attenuate the activity of viruses including retroviruses. Smap 29 has been demonstrated to have efficacy in reducing the activity of three retroviruses: namely HIV,  
15 HSV-1 and EIAV.

This invention thus encompasses methods to inhibit the microbial growth through the use of synthetic peptides demonstrated to have antimicrobial properties. It is contemplated that these peptides may be delivered into an environment in which bacteria  
20 are present or are likely to be present in order to control their growth and proliferation. It is further envisioned that such an environment would include a host organism. These embodiments, as well as others, are set forth in the following detailed description of the invention.

## 25 1. Peptide production

### A. Peptide Synthesis

The antimicrobial peptides envisioned in the instant invention may be chemically synthesized. An example of a method for chemical synthesis of such a peptide is as follows. Using the solid phase peptide synthesis method of Sheppard *et al.* (1981) an  
30 automated peptide synthesizer (Pharmacia LKB Biotechnology Co., LKB Biotynk 4170) adds N,N'-dicyclohexylcarbodiimide to amino acids whose amine functional groups are

protected by 9-fluorenylmethoxycarbonyl groups, producing anhydrides of the desired amino acid (Fmoc-amino acids). An Fmoc amino acid corresponding to the C-terminal amino acid of the desired peptide is affixed to Ultrosyn A resin (Pharmacia LKB Biotechnology Co.) through its carboxyl group, using dimethylaminopyridine as a catalyst. The resin is then washed with dimethylformamide containing iperidine resulting in the removal of the protective amine group of the C-terminal amino acid. A Fmoc-amino acid anhydride corresponding to the next residue in the peptide sequence is then added to the substrate and allowed to couple with the unprotected amino acid affixed to the resin. The protective amine group is subsequently removed from the second amino acid and the above process is repeated with additional residues added to the peptide in a like manner until the sequence is completed. After the peptide is completed, the protective groups, other than the acetoamidomethyl group are removed and the peptide is released from the resin with a solvent consisting of, for example, 94% (by weight) trifluoroacetic acid, 5% phenol, and 1% ethanol. The synthesized peptide is subsequently purified using high-performance liquid chromatography or other peptide purification technique discussed below.

In designing alternate peptide constructs with enhanced antimicrobial properties, substitutions may be used which modulate one or more properties of the molecule. Such variants typically contain the exchange of one amino acid for another at one or more sites within the peptide. For example, certain amino acids may be substituted for other amino acids in a peptide structure in order to enhance the interactive binding capacity of the structures. Since it is the interactive capacity and nature of a protein that defines that protein's biological functional activity, certain amino acid substitutions can be made in a protein sequence, and its underlying DNA coding sequence which potentially create a peptide with superior characteristics.

In making such changes, the hydropathic index of amino acids may be considered. The importance of the hydropathic amino acid index in conferring interactive biologic function on a protein is generally understood in the art (Kyte & Doolittle, 1982). It is accepted that the relative hydropathic character of the amino acid contributes to the

secondary structure of the resultant protein, which in turn defines the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens, and the like.

5        Each amino acid has been assigned a hydropathic index on the basis of their hydrophobicity and charge characteristics (Kyte & Doolittle, 1982), these are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5);  
10    glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

It is known in the art that certain amino acids may be substituted by other amino acids having a similar hydropathic index or score and still result in a protein with similar biological activity, *i.e.*, still obtain a biological functionally equivalent protein. In making  
15    such changes, the substitution of amino acids whose hydropathic indices are within  $\pm 2$  is preferred, those which are within  $\pm 1$  are particularly preferred, and those within  $\pm 0.5$  are even more particularly preferred.

It is also understood in the art that the substitution of like amino acids can be  
20    made effectively on the basis of hydrophilicity. U.S. Patent 4,554,101, incorporated herein by reference, states that the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a biological property of the protein. As detailed in U.S. Patent 4,554,101, the following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine  
25    (+3.0); aspartate (+3.0  $\pm$  1); glutamate (+3.0  $\pm$  1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5  $\pm$  1); alanine (-0.5); histidine \*-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); tryptophan (-3.4).

30        Amino acid substitutions are generally based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity,

charge, size, and the like but may nevertheless be made to highlight a particular property of the peptide. Exemplary substitutions that take various of the foregoing characteristics into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and  
5 valine, leucine and isoleucine.

#### B. *Fusion Proteins*

The antimicrobial peptides of the instant application may be combined with fusion partners to produce fusion proteins. It is envisioned that such constructs might  
10 include combinations of an antimicrobial peptide with a partner also exhibiting some level of antimicrobial activity. Such a construct generally has all or a substantial portion of the native molecule, linked at the N- or C-terminus, to all or a portion of a second polypeptide. For example, fusions typically employ leader sequences from other species to permit the recombinant expression of a protein in a heterologous host. Another useful  
15 fusion includes the addition of an immunologically active domain, such as an antibody epitope, to facilitate purification of the fusion protein. Inclusion of a cleavage site at or near the fusion junction will facilitate removal of the extraneous polypeptide after purification if such removal is desired. Other useful fusions include linking of functional domains, such as active sites from enzymes, glycosylation domains, cellular targeting  
20 signals or transmembrane regions.

It is envisioned that, to construct fusion proteins, the cDNA sequence encoding the antimicrobial peptide would be linked to the cDNA sequence encoding the desired fusion partner. The antimicrobial peptide sequences disclosed in this application allow  
25 for the deduction of encoding DNA. Such sequences may be prepared using conventional techniques, and used as probes to recover corresponding DNA's from genomic or cDNA libraries. Following cloning, such DNA's can then be incorporated in appropriate expression vectors and used to transform host cells (e.g., bacterial or mammalian cells), which can be cultured to form recombinant antimicrobial peptides.

30

As used in this application, the term "an isolated nucleic acid encoding a antimicrobial peptide refers to a nucleic acid molecule that has been isolated free of total cellular nucleic acid. The term "functionally equivalent codon" is used herein to refer to codons that encode the same amino acid, such as the six codons for arginine or serine (Table 1, below), and also refers to codons that encode biologically equivalent amino acids, as discussed in the following pages.



TABLE 1: CODONS

Amino Acids			Codons
Alanine	Ala	A	GCA GCC GCG GCU
Cysteine	Cys	C	UGC UGU
Aspartic acid	Asp	D	GAC GAU
Glutamic acid	Glu	E	GAA GAG
Phenylalanine	Phe	F	UUC UUU
Glycine	Gly	G	GGA GGC GGG GGU
Histidine	His	H	CAC CAU
Isoleucine	Ile	I	AUA AUC AUU
Lysine	Lys	K	AAA AAG
Leucine	Leu	L	UUA UUG CUA CUC CUG CUU
Methionine	Met	M	AUG
Asparagine	Asn	N	AAC AAU
Proline	Pro	P	CCA CCC CCG CCU
Glutamine	Gln	Q	CAA CAG
Arginine	Arg	R	AGA AGG CGA CGC CGG CGU
Serine	Ser	S	AGC AGU UCA UCC UCG UCU
Threonine	Thr	T	ACA ACC ACG ACU
Valine	Val	V	GUA GUC GUG GUU
Tryptophan	Trp	W	UGG
Tyrosine	Tyr	Y	UAC UAU

Allowing for the degeneracy of the genetic code, sequences that have at least about 50%, usually at least about 60%, more usually about 70%, most usually about 80%, preferably at least about 90% and most preferably about 95% of nucleotides that are identical to the nucleotides of an antimicrobial peptide gene will be sequences that encompassed by the present invention. Nucleic acid sequences of the present invention may also be functionally defined as sequences that are capable of hybridizing to a nucleic acid segment encoding an antimicrobial peptide.

The DNA segments of the present invention include those encoding biologically functional equivalent antimicrobial peptides, as described above. Functionally equivalent proteins or peptides may be created via the application of recombinant DNA technology, in which changes in the protein structure may be engineered, based on considerations of the properties of the amino acids being exchanged, or as a result of natural selection. Changes designed by man may be introduced through the application of site-directed mutagenesis techniques or may be introduced randomly and screened later for the desired function.

10 C. *Gene Therapy*

In other embodiments, it is envisioned that antimicrobial peptides may be utilized in gene therapy. Individuals who are immunodeficient due to disease, injury or genetic defect may be the subject of gene therapy in which the genes for antimicrobial peptides are incorporated into host cells. To facilitate gene therapy, the cDNA for antimicrobial peptides must be incorporated into an expression construct.

Expression requires that appropriate signals be provided in the vectors, and which include various regulatory elements, such as enhancers/promoters from both viral and mammalian sources that drive expression of the genes of interest in host cells. Elements designed to optimize messenger RNA stability and translatability in host cells also are defined. The conditions for the use of a number of dominant drug selection markers for establishing permanent, stable cell clones expressing the products are also provided, as is an element that links expression of the drug selection markers to expression of the polypeptide.

25

In general, plasmid vectors containing replicon and control sequences which are derived from species compatible with the host cell are used in connection with these hosts. The vector ordinarily carries a replication site, as well as marking sequences which are capable of providing phenotypic selection in transformed cells. For example, *E. coli* is often transformed using derivatives of pBR322, a plasmid derived from an *E. coli* species. pBR322 contains genes for ampicillin and tetracycline resistance and thus

30

provides easy means for identifying transformed cells. The pBR plasmid, or other microbial plasmid or phage must also contain, or be modified to contain, promoters which can be used by the microbial organism for expression of its own proteins.

5 In addition, phage vectors containing replicon and control sequences that are compatible with the host microorganism can be used as transforming vectors in connection with these hosts. For example, the phage lambda GEM<sup>TM</sup>-11 may be utilized in making a recombinant phage vector which can be used to transform host cells, such as *E. coli* LE392.

10

Further useful vectors include pIN vectors (Inouye *et al.*, 1985); and pGEX vectors, for use in generating glutathione S-transferase (GST) soluble fusion proteins for later purification and separation or cleavage. Other suitable fusion proteins are those with  $\beta$ -galactosidase, ubiquitin, the like.

15

Viral vectors are preferred eukaryotic expression systems. Included are adenoviruses, adeno-associated viruses, retroviruses, herpesviruses, poxviruses including vaccinia viruses and papilloma viruses including SV40.

20

#### (i) Regulatory Elements

Throughout this application, the term "expression construct" is meant to include any type of genetic construct containing a polynucleotide coding for a gene product in which part or all of the nucleic acid encoding sequence is capable of being transcribed. The transcript may be translated into a protein, but it need not be. In certain  
25 embodiments, expression includes both transcription of a gene and translation of mRNA into a gene product. In other embodiments, expression only includes transcription of the nucleic acid encoding a gene of interest.

In preferred embodiments, the nucleic acid encoding a gene product is under  
30 transcriptional control of a promoter. A "promoter" refers to a DNA sequence recognized by the synthetic machinery of the cell, or introduced synthetic machinery, required to

initiate the specific transcription of a gene. The phrase "under transcriptional control" means that the promoter is in the correct location and orientation in relation to the nucleic acid to control RNA polymerase initiation and expression of the gene.

5       The term eukaryotic promoter will be used here to refer to a group of transcriptional control modules that are clustered around the initiation site for RNA polymerase II. Much of the thinking about how promoters are organized derives from analyses of several viral promoters, including those for the HSV thymidine kinase (*tk*) and SV40 early transcription units. These studies, augmented by more recent work, have  
10 shown that promoters are composed of discrete functional modules, each consisting of approximately 7-20 bp of DNA, and containing one or more recognition sites for transcriptional activator or repressor proteins.

At least one module in each promoter functions to position the start site for RNA  
15 synthesis. The best known example of this is the TATA box, but in some promoters lacking a TATA box, such as the promoter for the mammalian terminal deoxynucleotidyl transferase gene and the promoter for the SV40 late genes, a discrete element overlying the start site itself helps to fix the place of initiation.

20       Additional promoter elements regulate the frequency of transcriptional initiation. Typically, these are located in the region 30-110 bp upstream of the start site, although a number of promoters have recently been shown to contain functional elements downstream of the start site as well. The spacing between promoter elements frequently is flexible, so that promoter function is preserved when elements are inverted or moved  
25 relative to one another. In the *tk* promoter, the spacing between promoter elements can be increased to 50 bp apart before activity begins to decline. Depending on the promoter, it appears that individual elements can function either co-operatively or independently to activate transcription.

30       The particular promoter employed to control the expression of a nucleic acid sequence of interest is not believed to be important, so long as it is capable of direction

the expression of the nucleic acid in the targeted cell. Thus, where a human cell is targeted, it is preferable to position the nucleic acid coding region adjacent to and under the control of a promoter that is capable of being expressed in a human cell. Generally speaking, such a promoter might include either a human or viral promoter.

5

In various embodiments, the human cytomegalovirus (CMV) immediate early gene promoter, the SV40 early promoter, the Rous sarcoma virus long terminal repeat, rat insulin promoter and glyceraldehyde-3-phosphate dehydrogenase can be used to obtain high-level expression of the coding sequence of interest. The use of other viral or  
10 mammalian cellular or bacterial phage promoters which are well-known in the art to achieve expression of a coding sequence of interest is contemplated as well, provided that the levels of expression are sufficient for a given purpose.

By employing a promoter with well-known properties, the level and pattern of  
15 expression of the protein of interest following transfection or transformation can be optimized. Further, selection of a promoter that is regulated in response to specific physiologic signals can permit inducible expression of the gene product.

Enhancers are genetic elements that increase transcription from a promoter  
20 located at a distant position on the same molecule of DNA. Enhancers are organized much like promoters. That is, they are composed of many individual elements, each of which binds to one or more transcriptional proteins.

The basic distinction between enhancers and promoters is operational. An  
25 enhancer region as a whole must be able to stimulate transcription at a distance; this need not be true of a promoter region or its component elements. On the other hand, a promoter must have one or more elements that direct initiation of RNA synthesis at a particular site and in a particular orientation, whereas enhancers lack these specificities. Promoters and enhancers are often overlapping and contiguous, often seeming to have a  
30 very similar modular organization.



Where a cDNA insert is employed, one will typically desire to include a polyadenylation signal to effect proper polyadenylation of the gene transcript. The nature of the polyadenylation signal is not believed to be crucial to the successful practice of the invention, and any such sequence may be employed such as human growth hormone and SV40 polyadenylation signals. Also contemplated as an element of the expression cassette is a terminator. These elements can serve to enhance message levels and to minimize read through from the cassette into other sequences.

(ii) Selectable Markers

In certain embodiments of the invention, the cells contain nucleic acid constructs of the present invention, a cell may be identified *in vitro* or *in vivo* by including a marker in the expression construct. Such markers would confer an identifiable change to the cell permitting easy identification of cells containing the expression construct. Usually the inclusion of a drug selection marker aids in cloning and in the selection of transformants, for example, genes that confer resistance to neomycin, puromycin, hygromycin, DHFR, GPT, zeocin and histidinol are useful selectable markers. Alternatively, enzymes such as herpes simplex virus thymidine kinase (*tk*) or chloramphenicol acetyltransferase (CAT) may be employed. Immunologic markers also can be employed. The selectable marker employed is not believed to be important, so long as it is capable of being expressed simultaneously with the nucleic acid encoding a gene product. Further examples of selectable markers are well known to one of skill in the art.

(iii) Multigene Constructs and IRES

In certain embodiments of the invention, the use of internal ribosome binding sites (IRES) elements are used to create multigene, or polycistronic, messages. IRES elements are able to bypass the ribosome scanning model of 5' methylated Cap dependent translation and begin translation at internal sites (Pelletier and Sonenberg, 1988). IRES elements from two members of the picornavirus family (polio and encephalomyocarditis) have been described (Pelletier and Sonenberg, 1988), as well an IRES from a mammalian message (Macejak and Sarnow, 1991). IRES elements can be linked to heterologous open reading frames. Multiple open reading frames can be transcribed together, each

separated by an IRES, creating polycistronic messages. By virtue of the IRES element, each open reading frame is accessible to ribosomes for efficient translation. Multiple genes can be efficiently expressed using a single promoter/enhancer to transcribe a single message.

5

Any heterologous open reading frame can be linked to IRES elements. This includes genes for secreted proteins, multi-subunit proteins, encoded by independent genes, intracellular or membrane-bound proteins and selectable markers. In this way, expression of several proteins can be simultaneously engineered into a cell with a single construct and a single selectable marker.

10

#### (iv) Host Cells and Delivery of Expression Vectors

Certain examples of prokaryotic hosts are *E. coli* strain RR1, *E. coli* LE392, *E. coli* B, *E. coli* X 1776 (ATCC No. 31537) as well as *E. coli* W3110 (F-, lambda-, prototrophic, ATCC No. 273325); bacilli such as *Bacillus subtilis*; and other enterobacteriaceae such as *Salmonella typhimurium*, *Serratia marcescens*, and various *Pseudomonas* species.

15

Primary mammalian cell cultures may be prepared in various ways. In order for the cells to be kept viable while *in vitro* and in contact with the expression construct, it is necessary to ensure that the cells maintain contact with the correct ratio of oxygen and carbon dioxide and nutrients but are protected from microbial contamination. Cell culture techniques are well documented and are disclosed herein by reference (Freshner, 1992).

20

There are a number of ways in which expression vectors may introduced into cells. In certain embodiments of the invention, the expression construct comprises a virus or engineered construct derived from a viral genome. The ability of certain viruses to enter cells via receptor-mediated endocytosis, to integrate into host cell genome and express viral genes stably and efficiently have made them attractive candidates for the transfer of foreign genes into mammalian cells (Ridgeway, 1988; Nicolas and Rubenstein, 1988; Baichwal and Sugden, 1986; Temin, 1986). The first viruses used as

25

30

gene vectors were DNA viruses including the papovaviruses (simian virus 40, bovine papilloma virus, and polyoma) (Ridgeway, 1988; Baichwal and Sugden, 1986) and adenoviruses (Ridgeway, 1988; Baichwal and Sugden, 1986). These have a relatively low capacity for foreign DNA sequences and have a restricted host spectrum. Furthermore, their oncogenic potential and cytopathic effects in permissive cells raise safety concerns. They can accommodate only up to 8 kb of foreign genetic material but can be readily introduced in a variety of cell lines and laboratory animals (Nicolas and Rubenstein, 1988; Temin, 1986).

One of the preferred methods for *in vivo* delivery involves the use of an adenovirus expression vector. "Adenovirus expression vector" is meant to include those constructs containing adenovirus sequences sufficient to (a) support packaging of the construct and (b) to express an antisense polynucleotide that has been cloned therein. In this context, expression does not require that the gene product be synthesized.

The expression vector comprises a genetically engineered form of adenovirus. Knowledge of the genetic organization of adenovirus, a 36 kb, linear, double-stranded DNA virus, allows substitution of large pieces of adenoviral DNA with foreign sequences up to 7 kb (Grunhaus & Horwitz, 1992). In contrast to retrovirus, the adenoviral infection of host cells does not result in chromosomal integration because adenoviral DNA can replicate in an episomal manner without potential genotoxicity. Also, adenoviruses are structurally stable, and no genome rearrangement has been detected after extensive amplification. Adenovirus can infect virtually all epithelial cells regardless of their cell cycle stage. So far, adenoviral infection appears to be linked only to mild disease such as acute respiratory disease in humans.

Adenovirus is particularly suitable for use as a gene transfer vector because of its mid-sized genome, ease of manipulation, high titer, wide target cell range and high infectivity. Both ends of the viral genome contain 100-200 base pair inverted repeats (ITRs), which are *cis* elements necessary for viral DNA replication and packaging. The early (E) and late (L) regions of the genome contain different transcription units that are

divided by the onset of viral DNA replication. The E1 region (E1A and E1B) encodes proteins responsible for the regulation of transcription of the viral genome and a few cellular genes. The expression of the E2 region (E2A and E2B) results in the synthesis of the proteins for viral DNA replication. These proteins are involved in DNA replication,  
5 late gene expression and host cell shut-off (Renan, 1990). The products of the late genes, including the majority of the viral capsid proteins, are expressed only after significant processing of a single primary transcript issued by the major late promoter (MLP). The MLP, (located at 16.8 m.u.) is particularly efficient during the late phase of infection, and all the mRNA's issued from this promoter possess a 5'-tripartite leader (TPL) sequence  
10 which makes them preferred mRNA's for translation.

In a current system, recombinant adenovirus is generated from homologous recombination between shuttle vector and provirus vector. Due to the possible recombination between two proviral vectors, wild-type adenovirus may be generated  
15 from this process. Therefore, it is critical to isolate a single clone of virus from an individual plaque and examine its genomic structure.

Generation and propagation of the current adenovirus vectors, which are replication deficient, depend on a unique helper cell line, designated 293, which was  
20 transformed from human embryonic kidney cells by Ad5 DNA fragments and constitutively expresses E1 proteins (Graham *et al.*, 1977). Since the E3 region is dispensable from the adenovirus genome (Jones and Shenk, 1978), the current adenovirus vectors, with the help of 293 cells, carry foreign DNA in either the E1, the D3 or both regions (Graham and Prevec, 1991). In nature, adenovirus can package approximately  
25 105% of the wild-type genome (Ghosh-Choudhury *et al.*, 1987), providing capacity for about 2 extra kb of DNA. Combined with the approximately 5.5 kb of DNA that is replaceable in the E1 and E3 regions, the maximum capacity of the current adenovirus vector is under 7.5 kb, or about 15% of the total length of the vector. More than 80% of the adenovirus viral genome remains in the vector backbone and is the source of vector-  
30 borne cytotoxicity. Also, the replication deficiency of the E1-deleted virus is incomplete.

For example, leakage of viral gene expression has been observed with the currently available vectors at high multiplicities of infection (MOI) (Mulligan, 1993).

Helper cell lines may be derived from human cells such as human embryonic  
5 kidney cells, muscle cells, hematopoietic cells or other human embryonic mesenchymal  
or epithelial cells. Alternatively, the helper cells may be derived from the cells of other  
mammalian species that are permissive for human adenovirus. Such cells include, *e.g.*,  
Vero cells or other monkey embryonic mesenchymal or epithelial cells. As stated above,  
the preferred helper cell line is 293.

10

Racher *et al.* (1995) discloses improved methods for culturing 293 cells and  
propagating adenovirus. In one format, natural cell aggregates are grown by inoculating  
individual cells into 1 liter siliconized spinner flasks (Techne, Cambridge, UK)  
containing 100-200 ml of medium. Following stirring at 40 rpm, the cell viability is  
15 estimated with trypan blue. In another format, Fibra-Cel microcarriers (Bibby Sterlin,  
Stone, UK) (5 g/l) is employed as follows. A cell inoculum, resuspended in 5 ml of  
medium, is added to the carrier (50 ml) in a 250 ml Erlenmeyer flask and left stationary,  
with occasional agitation, for 1 to 4 hours. The medium is then replaced with 50 ml of  
fresh medium and shaking initiated. For virus production, cells are allowed to grow to  
20 about 80% confluence, after which time the medium is replaced (to 25% of the final  
volume) and adenovirus added at an MOI of 0.05. Cultures are left stationary overnight,  
following which the volume is increased to 100% and shaking commenced for another 72  
hours.

25

Other than the requirement that the adenovirus vector be replication defective, or  
at least conditionally defective, the nature of the adenovirus vector is not believed to be  
crucial to the successful practice of the invention. The adenovirus may be of any of the  
42 different known serotypes or subgroups A-F. Adenovirus type 5 of subgroup C is the  
preferred starting material in order to obtain the conditional replication-defective  
30 adenovirus vector for use in the present invention. This is because Adenovirus type 5 is a  
human adenovirus about which a great deal of biochemical and genetic information is



known, and it has historically been used for most constructions employing adenovirus as a vector.

As stated above, the typical vector according to the present invention is replication defective and will not have an adenovirus E1 region. Thus, it will be most convenient to introduce the polynucleotide encoding the gene of interest at the position from which the E1-coding sequences have been removed. However, the position of insertion of the construct within the adenovirus sequences is not critical to the invention. The polynucleotide encoding the gene of interest may also be inserted in lieu of the deleted E3 region in E3 replacement vectors as described by Karlsson *et al.*, (1986) or in the E4 region where a helper cell line or helper virus complements the E4 defect.

Adenovirus is easy to grow and manipulate and exhibits broad host range *in vitro* and *in vivo*. This group of viruses can be obtained in high titers, *e.g.*,  $10^9$ - $10^{11}$  plaque-forming units per ml, and they are highly infective. The life cycle of adenovirus does not require integration into the host cell genome. The foreign genes delivered by adenovirus vectors are episomal and, therefore, have low genotoxicity to host cells. No side effects have been reported in studies of vaccination with wild-type adenovirus (Top *et al.*, 1971), demonstrating their safety and therapeutic potential as *in vivo* gene transfer vectors.

Adenovirus vectors have been used in eukaryotic gene expression (Levrero *et al.*, 1991; Gomez-Foix *et al.*, 1992) and vaccine development (Grunhaus and Horwitz, 1992; Graham and Prevec, 1992). Recently, animal studies suggested that recombinant adenovirus could be used for gene therapy (Stratford-Perricaudet and Perricaudet, 1991; Stratford-Perricaudet *et al.*, 1990; Rich *et al.*, 1993). Studies in administering recombinant adenovirus to different tissues include trachea instillation (Rosenfeld *et al.*, 1991; Rosenfeld *et al.*, 1992), muscle injection (Ragot *et al.*, 1993), peripheral intravenous injections (Herz and Gerard, 1993) and stereotactic inoculation into the brain (Le Gal La Salle *et al.*, 1993).

The retroviruses are a group of single-stranded RNA viruses characterized by an ability to convert their RNA to double-stranded DNA in infected cells by a process of reverse-transcription (Coffin, 1990). The resulting DNA then stably integrates into cellular chromosomes as a provirus and directs synthesis of viral proteins. The  
5 integration results in the retention of the viral gene sequences in the recipient cell and its descendants. The retroviral genome contains three genes, gag, pol, and env that code for capsid proteins, polymerase enzyme, and envelope components, respectively. A sequence found upstream from the gag gene contains a signal for packaging of the genome into virions. Two long terminal repeat (LTR) sequences are present at the 5' and  
10 3' ends of the viral genome. These contain strong promoter and enhancer sequences and are also required for integration in the host cell genome (Coffin, 1990).

In order to construct a retroviral vector, a nucleic acid encoding a gene of interest is inserted into the viral genome in the place of certain viral sequences to produce a virus  
15 that is replication-defective. In order to produce virions, a packaging cell line containing the gag, pol, and env genes but without the LTR and packaging components is constructed (Mann *et al.*, 1983). When a recombinant plasmid containing a cDNA, together with the retroviral LTR and packaging sequences is introduced into this cell line (by calcium phosphate precipitation for example), the packaging sequence allows the  
20 RNA transcript of the recombinant plasmid to be packaged into viral particles, which are then secreted into the culture media (Nicolas and Rubenstein, 1988; Temin, 1986; Mann *et al.*, 1983). The media containing the recombinant retroviruses is then collected, optionally concentrated, and used for gene transfer. Retroviral vectors are able to infect a broad variety of cell types. However, integration and stable expression require the  
25 division of host cells (Paskind *et al.*, 1975).

A novel approach designed to allow specific targeting of retrovirus vectors was recently developed based on the chemical modification of a retrovirus by the chemical addition of lactose residues to the viral envelope. This modification could permit the  
30 specific infection of hepatocytes via sialoglycoprotein receptors.

A different approach to targeting of recombinant retroviruses was designed in which biotinylated antibodies against a retroviral envelope protein and against a specific cell receptor were used. The antibodies were coupled via the biotin components by using streptavidin (Roux *et al.*, 1989). Using antibodies against major histocompatibility  
5 complex class I and class II antigens, they demonstrated the infection of a variety of human cells that bore those surface antigens with an ecotropic virus *in vitro* (Roux *et al.*, 1989).

There are certain limitations to the use of retrovirus vectors in all aspects of the  
10 present invention. For example, retrovirus vectors usually integrate into random sites in the cell genome. This can lead to insertional mutagenesis through the interruption of host genes or through the insertion of viral regulatory sequences that can interfere with the function of flanking genes (Varmus *et al.*, 1981). Another concern with the use of defective retrovirus vectors is the potential appearance of wild-type replication-  
15 competent virus in the packaging cells. This can result from recombination events in which the intact- sequence from the recombinant virus inserts upstream from the gag, pol, env sequence integrated in the host cell genome. However, new packaging cell lines are now available that should greatly decrease the likelihood of recombination (Markowitz *et al.*, 1988; Hersdorffer *et al.*, 1990).

20

Other viral vectors may be employed as expression constructs in the present invention. Vectors derived from viruses such as vaccinia virus (Ridgeway, 1988; Baichwal and Sugden, 1986; Coupar *et al.*, 1988) adeno-associated virus (AAV) (Ridgeway, 1988; Baichwal and Sugden, 1986; Hermonat and Muzycska, 1984) and  
25 herpesviruses may be employed. They offer several attractive features for various mammalian cells (Friedmann, 1989; Ridgeway, 1988; Baichwal and Sugden, 1986; Coupar *et al.*, 1988; Horwich *et al.*, 1990).

With the recent recognition of defective hepatitis B viruses, new insight was  
30 gained into the structure-function relationship of different viral sequences. *In vitro* studies showed that the virus could retain the ability for helper-dependent packaging and

reverse transcription despite the deletion of up to 80% of its genome (Horwich *et al.*, 1990). This suggested that large portions of the genome could be replaced with foreign genetic material. The hepatotropism and persistence (integration) were particularly attractive properties for liver-directed gene transfer. Chang *et al.*, recently introduced the  
5 chloramphenicol acetyltransferase (CAT) gene into duck hepatitis B virus genome in the place of the polymerase, surface, and pre-surface coding sequences. It was co-transfected with wild-type virus into an avian hepatoma cell line. Culture media containing high titers of the recombinant virus were used to infect primary duckling hepatocytes. Stable CAT gene expression was detected for at least 24 days after transfection (Chang *et al.*,  
10 1991).

In order to effect expression of gene constructs, the expression construct must be delivered into a cell. This delivery may be accomplished *in vitro*, as in laboratory procedures for transforming cells lines, or *in vivo* or *ex vivo*, as in the treatment of certain  
15 disease states. One mechanism for delivery is via viral infection where the expression construct is encapsidated in an infectious viral particle.

Several non-viral methods for the transfer of expression constructs into cultured mammalian cells also are contemplated by the present invention. These include calcium  
20 phosphate precipitation (Graham and Van Der Eb, 1973; Chen and Okayama, 1987; Rippe *et al.*, 1990) DEAE-dextran (Gopal, 1985), electroporation (Tur-Kaspa *et al.*, 1986; Potter *et al.*, 1984), direct microinjection (Harland and Weintraub, 1985), DNA-loaded liposomes (Nicolau and Sene, 1982; Fraley *et al.*, 1979) and lipofectamine-DNA complexes, cell sonication (Fechheimer *et al.*, 1987), gene bombardment using high  
25 velocity microprojectiles (Yang *et al.*, 1990), and receptor-mediated transfection (Wu and Wu, 1987; Wu and Wu, 1988). Some of these techniques may be successfully adapted for *in vivo* or *ex vivo* use.

Once the expression construct has been delivered into the cell the nucleic acid  
30 encoding the gene of interest may be positioned and expressed at different sites. In certain embodiments, the nucleic acid encoding the gene may be stably integrated into the

genome of the cell. This integration may be in the cognate location and orientation via homologous recombination (gene replacement) or it may be integrated in a random, non-specific location (gene augmentation). In yet further embodiments, the nucleic acid may be stably maintained in the cell as a separate, episomal segment of DNA. Such nucleic acid segments or "episomes" encode sequences sufficient to permit maintenance and replication independent of or in synchronization with the host cell cycle. How the expression construct is delivered to a cell and where in the cell the nucleic acid remains is dependent on the type of expression construct employed.

10 In yet another embodiment of the invention, the expression construct may simply consist of naked recombinant DNA or plasmids. Transfer of the construct may be performed by any of the methods mentioned above which physically or chemically permeabilize the cell membrane. This is particularly applicable for transfer *in vitro* but it may be applied to *in vivo* use as well. Dubensky *et al.* (1984) successfully injected polyomavirus DNA in the form of calcium phosphate precipitates into liver and spleen of adult and newborn mice demonstrating active viral replication and acute infection. Benvenisty and Neshif (1986) also demonstrated that direct intraperitoneal injection of calcium phosphate-precipitated plasmids results in expression of the transfected genes. It is envisioned that DNA encoding a gene of interest may also be transferred in a similar manner *in vivo* and express the gene product.

25 In still another embodiment of the invention for transferring a naked DNA expression construct into cells may involve particle bombardment. This method depends on the ability to accelerate DNA-coated microprojectiles to a high velocity allowing them to pierce cell membranes and enter cells without killing them (Klein *et al.*, 1987). Several devices for accelerating small particles have been developed. One such device relies on a high voltage discharge to generate an electrical current, which in turn provides the motive force (Yang *et al.*, 1990). The microprojectiles used have consisted of biologically inert substances such as tungsten or gold beads.

30



Selected organs including the liver, skin, and muscle tissue of rats and mice have been bombarded *in vivo* (Yang *et al.*, 1990; Zelenin *et al.*, 1991). This may require surgical exposure of the tissue or cells, to eliminate any intervening tissue between the gun and the target organ, *i.e.*, *ex vivo* treatment. Again, DNA encoding a particular gene  
5 may be delivered via this method and still be incorporated by the present invention.

In a further embodiment of the invention, the expression construct may be entrapped in a liposome. Liposomes are vesicular structures characterized by a phospholipid bilayer membrane and an inner aqueous medium. Multilamellar liposomes  
10 have multiple lipid layers separated by aqueous medium. They form spontaneously when phospholipids are suspended in an excess of aqueous solution. The lipid components undergo self-rearrangement before the formation of closed structures and entrap water and dissolved solutes between the lipid bilayers (Ghosh and Bachhawat, 1991). Also contemplated are lipofectamine-DNA complexes.

15

Liposome-mediated nucleic acid delivery and expression of foreign DNA *in vitro* has been very successful. Wong *et al.*, (1980) demonstrated the feasibility of liposome-mediated delivery and expression of foreign DNA in cultured chick embryo, HeLa and hepatoma cells. Nicolau *et al.*, (1987) accomplished successful liposome-mediated gene  
20 transfer in rats after intravenous injection.

In certain embodiments of the invention, the liposome may be complexed with a hemagglutinating virus (HVJ). This has been shown to facilitate fusion with the cell membrane and promote cell entry of liposome-encapsulated DNA (Kaneda *et al.*, 1989).  
25 In other embodiments, the liposome may be complexed or employed in conjunction with nuclear non-histone chromosomal proteins (HMG-1) (Kato *et al.*, 1991). In yet further embodiments, the liposome may be complexed or employed in conjunction with both HVJ and HMG-1. In that such expression constructs have been successfully employed in transfer and expression of nucleic acid *in vitro* and *in vivo*, then they are applicable for  
30 the present invention. Where a bacterial promoter is employed in the DNA construct, it also will be desirable to include within the liposome an appropriate bacterial polymerase.



Other expression constructs which can be employed to deliver a nucleic acid encoding a particular gene into cells are receptor-mediated delivery vehicles. These take advantage of the selective uptake of macromolecules by receptor-mediated endocytosis in almost all eukaryotic cells. Because of the cell type-specific distribution of various receptors, the delivery can be highly specific (Wu and Wu, 1993).

Receptor-mediated gene targeting vehicles generally consist of two components: a cell receptor-specific ligand and a DNA-binding agent. Several ligands have been used for receptor-mediated gene transfer. The most extensively characterized ligands are asialoorosomucoid (ASOR) (Wu and Wu, 1987) and transferrin (Wagner *et al.*, 1990). Recently, a synthetic neoglycoprotein, which recognizes the same receptor as ASOR, has been used as a gene delivery vehicle (Ferkol *et al.*, 1993; Perales *et al.*, 1994) and epidermal growth factor (EGF) has also been used to deliver genes to squamous carcinoma cells (Myers, EPO 0273085).

In other embodiments, the delivery vehicle may comprise a ligand and a liposome. For example, Nicolau *et al.*, (1987) employed lactosyl-ceramide, a galactose-terminal asialganglioside, incorporated into liposomes and observed an increase in the uptake of the insulin gene by hepatocytes. Thus, it is feasible that a nucleic acid encoding a particular gene also may be specifically delivered into a cell type such as lung, epithelial or tumor cells, by any number of receptor-ligand systems with or without liposomes. For example, epidermal growth factor (EGF) may be used as the receptor for mediated delivery of a nucleic acid encoding a gene in many tumor cells that exhibit upregulation of EGF receptor. Mannose can be used to target the mannose receptor on liver cells. Also, antibodies to CD5 (CLL), CD22 (lymphoma), CD25 (T-cell leukemia) and MAA (melanoma) can similarly be used as targeting moieties.

In certain embodiments, gene transfer may more easily be performed under *ex vivo* conditions. *Ex vivo* gene therapy refers to the isolation of cells from an animal, the delivery of a nucleic acid into the cells *in vitro*, and then the return of the modified cells

back into an animal. This may involve the surgical removal of tissue/organs from an animal or the primary culture of cells and tissues.

#### D. *Preparations*

5 It is envisioned that the antimicrobial peptides and any second agents that might be delivered may be formulated and administered in any pharmacologically acceptable vehicle, such as parenteral, topical, aerosol, liposomal, nasal or ophthalmic preparations, with formulations designed for oral administration being currently preferred due to their ease of use. It is further envisioned that formulations antimicrobial peptides and any  
10 second agents that might be delivered may be formulated and administered in a manner that does not require that the be coupled with a pharmaceutically acceptable carrier. In those situations, it would be clear to one of ordinary skill in the art the types of diluents that would be proper for the proposed use of the peptides and any secondary agents required. Although further purification following synthesis may be desired, it is not  
15 necessarily required for use.

#### E. *Protein Purification*

Peptide purification techniques are well known to those of skill in the art. These techniques involve, at one level, the crude fractionation of the cellular milieu to  
20 polypeptide and non-polypeptide fractions. Having separated the polypeptide from other proteins, the polypeptide of interest may be further purified using chromatographic, immunologic and electrophoretic techniques to achieve partial or complete purification (or purification to homogeneity). Analytical methods particularly suited to the preparation of a pure peptide are ion-exchange chromatography, exclusion  
25 chromatography; polyacrylamide gel electrophoresis; isoelectric focusing. A particularly efficient method of purifying peptides is fast protein liquid chromatography or HPLC.

Certain aspects of the present invention concern the purification, and in particular embodiments, the substantial purification, of an encoded peptide. The term "purified  
30 peptide" as used herein, is intended to refer to a composition, isolatable from other components, wherein the peptide is purified to any degree relative to its naturally-

obtainable state. A purified peptide therefore also refers to a peptide, free from the environment in which it may naturally occur.

Generally, "purified" will refer to a peptide composition that has been subjected  
5 to fractionation to remove various other components, and which composition substantially retains its expressed biological activity. Where the term "substantially purified" is used, this designation will refer to a composition in which the protein or peptide forms the major component of the composition, such as constituting about 50%, about 60%, about 70%, about 80%, about 90%, about 95% or more peptides in the  
10 composition. The term "purified to homogeneity" is used to mean that the composition has been purified such that there is single protein species based on the particular test of purity employed for example SDS-PAGE or HPLC.

Various methods for quantifying the degree of purification of the peptide will be  
15 known to those of skill in the art in light of the present disclosure. These include, for example, assessing the amount of peptides within a fraction by SDS/PAGE analysis.

There is no general requirement that the peptide always be provided in their most purified state. Indeed, it is contemplated that less substantially purified products will  
20 have utility in certain embodiments. Partial purification may be accomplished by using fewer purification steps in combination, or by utilizing different forms of the same general purification scheme. For example, it is appreciated that a cation-exchange column chromatography performed utilizing an HPLC apparatus will generally result in a greater "-fold" purification than the same technique utilizing a low pressure  
25 chromatography system. Methods exhibiting a lower degree of relative purification may have advantages in total recovery of protein product, or in maintaining the activity of an expressed protein.

It is known that the migration of a peptide can vary, sometimes significantly, with  
30 different conditions of SDS/PAGE (Capaldi *et al.*, 1977). It will therefore be appreciated

that under differing electrophoresis conditions, the apparent molecular weights of purified or partially purified expression products may vary.

High Performance Liquid Chromatography (HPLC) is characterized by a very rapid separation with extraordinary resolution of peaks. This is achieved by the use of very fine particles and high pressure to maintain an adequate flow rate. Separation can be accomplished in a matter of minutes, or at most an hour. Moreover, only a very small volume of the sample is needed because the particles are so small and close-packed that the void volume is a very small fraction of the bed volume. Also, the concentration of the sample need not be very great because the bands are so narrow that there is very little dilution of the sample.

Affinity Chromatography is a chromatographic procedure that relies on the specific affinity between a substance to be isolated and a molecule that it can specifically bind to. This is a receptor-ligand type interaction. The column material is synthesized by covalently coupling one of the binding partners to an insoluble matrix. The column material is then able to specifically adsorb the substance from the solution. Elution occurs by changing the conditions to those in which binding will not occur (alter pH, ionic strength, temperature, *etc.*).

20

The matrix should be a substance that itself does not adsorb molecules to any significant extent and that has a broad range of chemical, physical and thermal stability. The ligand should be coupled in such a way as to not affect its binding properties. The ligand should also provide relatively tight binding. And it should be possible to elute the substance without destroying the sample or the ligand. One of the most common forms of affinity chromatography is immunoaffinity chromatography. The generation of antibodies that would be suitable for use in accord with the present invention is discussed below.

25

## 2. Therapeutic Uses

This invention encompasses methods to reduce antimicrobial resistance, caused by any of the seven mechanisms described by Davies (1986) above, using an antimicrobial peptide and one or more antimicrobial agents or antibiotics. A list of  
5 bacterial strains that have developed antibiotic resistance by one or more of these mechanisms is listed in Table 2 (Lorian, 1991).

The peptides have broad spectrum antimicrobial properties effective against both Gram-positive and Gram-negative strains of bacteria and have been demonstrated to kill  
10 strains previously deemed multiply drug resistant. An attractive feature of these peptides is their tolerance for high salt concentrations. The peptides maintain activity in physiological salt solutions and thus are envisioned to be effective in the treatment lung diseases such as cystic fibrosis.

15 The bacteriocidal properties of the antimicrobial peptides disclosed in combination with their stability and insensitivity to high salt concentrations allow them to be included in formulations to inhibit microbial growth and proliferation. The purified antimicrobial peptide may be used without further modifications or it may be diluted in a pharmaceutically acceptable carrier. Because of the stability of the peptides it is  
20 contemplated that the invention may be administered to humans or animals, included in food preparations, pharmaceutical preparations, medicinal and pharmaceutical products, cosmetic products, hygienic products, cleaning products and cleaning agents, as well as any material to which the peptides could be sprayed on or adhered to wherein the inhibition of microbial growth on such a material is desired.

25

The proper dosage of an antimicrobial peptide necessary to prevent microbial growth and proliferation depends upon a number of factors including the types of bacteria that might be present, the environment into which the peptide is being introduced, and the time that the peptide is envisioned to remain in a given area.

30

It is further contemplated that the antimicrobial peptides of the invention may be used in combination with or to enhance the activity of other antimicrobial agents or antibiotics. Combinations of the peptide with other agents may be useful to allow antibiotics to be used at lower doses due to toxicity concerns, to enhance the activity of  
5 antibiotics whose efficacy has been reduced or to effectuate a synergism between the components such that the combination is more effective than the sum of the efficacy of either component independently. Antibiotics which may be combined with an antimicrobial peptide in combination therapy include but are not limited to penicillin, ampicillin, amoxycillin, vancomycin, cycloserine, bacitracin, cephalosporin, imipenem,  
10 colistin, methicillin, streptomycin, kanamycin, tobramycin, gentamicin, tetracycline, chlortetracycline, doxycycline, chloramphenicol, lincomycin, clindamycin, erythromycin, oleandomycin, polymyxin nalidixic acid, rifamycin, rifampicin, gantrisin, trimethoprim, isoniazid, paraaminosalicylic acid, and ethambutol. Table 3 (Reese and Betts, 1993), lists the antibiotics generally preferred for use against a given pathogenic bacterium. It is  
15 contemplated that the effectiveness of all the antibiotics listed in Table 3 will be increased upon combination with an antimicrobial peptide. Table 4 (Reese and Betts, 1993), itemizes the common pathogenic bacteria that are implicated in focal infections. The present invention is thus contemplated for use against all such infections.



**TABLE 2:**  
**MECHANISMS OF RESISTANCE TO ANTIMICROBIAL AGENTS**

Antimicrobial Agent	Mechanisms Causing Resistance	Examples of Organisms
Aminoglycosides	Modifying enzymes: acetyltransferases, adenylyl- transferases (nucleotidyl- transferases), phosphotransferases Ribosomal resistance (streptomycin, spectinomycin)	Enterobacteriaceae, <i>P. aeruginosa</i> , <i>S. aureus</i> , <i>E. faecalis</i>
	Inadequate drug transport	<i>E. faecalis</i> , Enterobacteriaceae, <i>M. tuberculosis</i> , <i>P. aeruginosa</i> <i>E. faecalis</i> , <i>P. aeruginosa</i> , anaerobes
$\beta$ -Lactams	Enzymatic inactivation	<i>S. aureus</i> , <i>E. faecalis</i> , Enterobacteriaceae, <i>P. aeruginosa</i> , <i>Neisseria</i> spp., <i>H. influenzae</i>
	Low affinity PBPs	<i>S. pneumoniae</i> , <i>N. gonorrhoeae</i> , <i>S. aureus</i> ,
	Lack of penetration through outer membrane	<i>P. aeruginosa</i> <i>P. aeruginosa</i> , Enterobacteriaceae
Chloramphenicol	Acetylation	Enterobacteriaceae, <i>S. aureus</i> , streptococci, <i>Bacteroides uniformis</i>
	Lack of penetration	<i>P. aeruginosa</i>
Clindamycin, erythromycin, lincomycin	Ribosomal resistance due to methylation of rRNA Inactivation by esterase Decreased penetration	Streptococci, <i>E. faecalis</i> , Enterobacteriaceae Enterobacteriaceae <i>S. hominis</i>
Fluoroquinolones	Decreased uptake	Enterobacteriaceae, <i>P. aeruginosa</i> ,
	Altered target site (DNA gyrase)	staphylococci Enterobacteriaceae, <i>P. aeruginosa</i>

Antimicrobial Agent	Mechanisms Causing Resistance	Examples of Organisms
Lincomycin	Inactivation	<i>S. aureus</i>
Sulfonamides	Synthesis of an altered or alternative target site (dihydropteroate synthetase) Lack of penetration Overproduction of PABA	Enterobacteriaceae, <i>Neisseria</i> spp., <i>P. aeruginosa</i> Anaerobes <i>Neisseria</i> , <i>S. aureus</i>
Tetracycline	Drug efflux  Protection of ribosome from tetracycline  Inactivation	Enterobacteriaceae, staphylococci, streptococci Streptococci, <i>E. faecalis</i> , <i>Neisseria</i> spp., <i>Mycoplasma</i> spp. Cryptic gene found in <i>B. fragilis</i> , expressed resistance in <i>E. coli</i>
Trimethoprim	Synthesis of an altered or alternative target site (dihydrofolate reductase) Lack of penetration Ability to use alternative pathway Overproduction of dihydrofolate reductase	Enterobacteriaceae, <i>V. cholerae</i> , staphylococci  <i>P. aeruginosa</i> Enterococci <i>H. influenzae</i>
Vancomycin	?  ?Blocking of target site	Pediococci, <i>Leuconostoc</i> spp. (intrinsic) Enterococci (acquired)

**TABLE 3:**  
**ANTIBIOTICS OF CHOICE FOR COMMON PATHOGENS**

Pathogen	Antibiotic of First Choice <sup>a</sup>	Alternative Agents <sup>a</sup>
<b>Gram-positive cocci</b>		
<i>Staphylococcus aureus</i> or <i>S. epidermidis</i> Non-penicillinase-producing	Penicillin	A first-generation cephalosporin, vancomycin, imipenem, or clindamycin; a fluoroquinolone <sup>b</sup>
Penicillinase-producing	Penicillinase-resistant penicillin ( <i>e.g.</i> , oxacillin or nafcillin)	A first-generation cephalosporin, vancomycin, clindamycin, imipenem, amoxicillin-clavulanic acid, ticarcillin-clavulanic acid, ampicillin-sulbactam; a fluoroquinolone <sup>b</sup>
Methicillin-resistant	Vancomycin with or without gentamicin and/or rifampin	TMP-SMZ, minocycline
Streptococci Group A, C, G	Penicillin	A cephalosporin <sup>a</sup> , vancomycin, erythromycin; clarithromycin; azithromycin; clindamycin
Group B	Penicillin (or ampicillin)	A cephalosporin <sup>a</sup> , vancomycin, or erythromycin
Enterococcus Endocarditis or other serious infection	Penicillin (or ampicillin) with gentamicin	Vancomycin with gentamicin
Uncomplicated urinary tract infection	Ampicillin or amoxicillin	A fluoroquinolone, nitrofurantoin
Viridans group	Penicillin G (with or without gentamicin)	A cephalosporin <sup>a</sup> , vancomycin
<i>S. bovis</i>	Penicillin G	A cephalosporin <sup>a</sup> , vancomycin
<i>S. pneumoniae</i>	Penicillin G	A cephalosporin <sup>a</sup> , erythromycin, chloramphenicol, vancomycin
<b>Gram-negative cocci</b>		

<i>Neisseria gonorrhoeae</i>	Ceftriaxone	Spectinomycin, a fluoroquinolone, cefoxitin, cefixime, cefotaxime (see Appendix E)
<i>N. meningitidis</i>	Penicillin G	Third-generation cephalosporin, chloramphenicol
<i>Moraxella (Branhamella) catarrhalis</i>	TMP-SMZ	Amoxicillin-clavulanic acid; an erythromycin; clarithromycin azithromycin, cefuroxime, cefixime, third-generation cephalosporin, tetracycline
<b>Gram-positive bacilli</b>		
<i>Clostridium perfringens</i> (and <i>Clostridium</i> sp.)	Penicillin G	Chloramphenicol, metronidazole, or clindamycin
<i>Listeria monocytogenes</i>	Ampicillin with or without gentamicin	TMP-SMZ
<b>Gram-negative bacilli</b>		
<i>Acinetobacter</i>	Imipenem	Tobramycin, gentamicin, or amikacin, usually with ticarcillin or piperacillin (or similar agent); TMP-SMZ
<i>Aeromonas hydrophila</i>	TMP-SMZ	Gentamicin, tobramycin; imipenem; a fluoroquinolone
<i>Bacteroides Bacteroides</i> sp. (oropharyngeal) <i>B. fragilis</i> strains (gastrointestinal strains)	Penicillin G  Metronidazole	Clindamycin, cefoxitin, metronidazole, chloramphenicol, cefotetan, ampicillin-sulbactam Clindamycin; ampicillin-sulbactam; imipenem; cefoxitin <sup>c</sup> ; cefotetan <sup>c</sup> ; ticarcillin-clavulanic acid; piperacillin <sup>c</sup> ; chloramphenicol; cefmetazole <sup>c</sup>
<i>Campylobacter</i>	A fluoroquinolone (adults)	A tetracycline, gentamicin

<i>fetus, jejuni</i>	or an erythromycin	
<i>Enterobacter</i> sp.	Imipenem	An aminoglycoside and piperacillin or ticarcillin or mezlocillin; a third-generation cephalosporin <sup>d</sup> ; TMP-SMZ; aztreonam; a fluoroquinolone
<i>Escherichia coli</i> Uncomplicated urinary tract infection Recurrent or systemic infection	TMP-SMZ A cephalosporin <sup>e</sup>	A cephalosporin or a fluoroquinolone  Ampicillin with or without an aminoglycoside, TMP-SMZ, oral fluoroquinolones useful in recurrent infections, ampicillin-sulbactam, ticarcillin-clavulanic acid, aztreonam
<i>Haemophilus influenzae</i> (cocci bacillary) Life-threatening infections Upper respiratory infections and bronchitis	Cefotaxime or ceftriaxone TMP-SMZ	Chloramphenicol; cefuroxime for pneumonia) Ampicillin or amoxicillin; cefuroxime; a sulfonamide with or without an erythromycin; cefuroxime-axetil; third-generation cephalosporin, amoxicillin-clavulanic acid, cefaclor, tetracycline; clarithromycin; azithromycin
<i>Klebsiella pneumoniae</i>	A cephalosporin <sup>e</sup>	An aminoglycoside, imipenem, TMP-SMZ, ticarcillin-clavulanic acid, ampicillin-sulbactam, aztreonam, a fluoroquinolone; amoxicillin-clavulanic acid
<i>Legionella</i> spp.	Erythromycin with rifampin	TMP-SMZ; clarithromycin; azithromycin; ciprofloxacin
<i>Pasteurella multocida</i>	Penicillin G	Tetracycline, cefuroxime, amoxicillin-clavulanic acid,

		ampicillin-sulbactam
<i>Proteus</i> sp.	Cefotaxime, ceftizoxime, or ceftriaxone <sup>f</sup>	An aminoglycoside; ticarcillin or piperacillin or mezlocillin; TMP-SMZ; amoxicillin-clavulanic acid; ticarcillin-clavulanic acid, ampicillin-sulbactam; a fluoroquinolone; aztreonam; imipenem
<i>Providencia stuartii</i>	Cefotaxime, ceftizoxime, or ceftriaxone <sup>f</sup>	Imipenem; an aminoglycoside often combined with ticarcillin or piperacillin or similar agent; ticarcillin-clavulanic acid; TMP-SMZ, a fluoroquinolone; aztreonam
<i>Pseudomonas aeruginosa</i> (nonurinary tract infection)  (urinary tract infections)	Gentamicin or tobramycin or amikacin (combined with ticarcillin, piperacillin, etc. for serious infections) Ciprofloxacin	An aminoglycoside and ceftazidime; imipenem, or aztreonam plus an aminoglycoside; ciprofloxacin  Carbenicillin; ticarcillin, piperacillin, or mezlocillin; ceftazidime; imipenem; aztreonam; an aminoglycoside
<i>Pseudomonas cepacia</i>	TMP-SMZ	Ceftazidime, chloramphenicol
<i>Salmonella typhi</i>  Other species	Ceftriaxone  Cefotaxime or ceftriaxone	Ampicillin, amoxicillin, TMP-SMZ, chloramphenicol; a fluoroquinolone Ampicillin or amoxicillin, TMP-SMZ, chloramphenicol; a fluoroquinolone
<i>Serratia</i>	Cefotaxime, ceftizoxime, or ceftriaxone <sup>f</sup>	Gentamicin or amikacin; imipenem; TMP-SMZ; ticarcillin, piperacillin, or mezlocillin; aztreonam; a



		fluoroquinolone
<i>Shigella</i>	A fluoroquinolone	TMP-SMZ; ceftriaxone; ampicillin
<i>Vibrio cholerae</i> (chlorea)	A tetracycline	TMP-SMZ; a fluoroquinolone
<i>Vibrio vulnificus</i>	A tetracycline	Cefotaxime
<i>Xanthomonas</i> ( <i>Pseudomonas</i> ) <i>maltophilia</i>	TMP-SMZ	Minocycline, ceftazidime, a fluoroquinolone
<i>Yersinia enterocolitica</i>	TMP-SMZ	A fluoroquinolone; an aminoglycoside; cefotaxime or ceftizoxime
<i>Yersinia pestis</i> (plague)	Streptomycin	A tetracycline; chloramphenicol; gentamicin

Key: TMP-SMZ = trimethoprim-sulfamethoxazole.

<sup>a</sup> Choice presumes susceptibility studies indicate that the pathogen is susceptible to the agent.

5 <sup>b</sup>The experience with fluoroquinolone use in staphylococcal infections is relatively limited.

The fluoroquinolones should be used only in adults.

<sup>c</sup>Up to 15-20% of strains may be resistant.

<sup>d</sup>*Enterobacter spp.* may develop resistance to the cephalosporins.

10 <sup>e</sup>Specific choice will depend on susceptibility studies. Third-generation cephalosporins may be exquisitely active against many Gram-negative bacilli (e.g., *E. coli*, *Klebsiella* sp.). In some geographic areas, 20-25% of community-acquired *E. coli* infections may be resistant to ampicillin (amoxicillin).

15 <sup>f</sup>In severely ill patients, this is often combined with an aminoglycoside while awaiting susceptibility data.

**TABLE 4:**  
**COMMON PATHOGENS IN FOCAL INFECTIONS**

Presumed location of Infection	Common pathogens	Gram stain Characteristics of exudate-if available
Urinary tract infections	Community-acquired: <i>Escherichia coli</i> Recurrent or nosocomial: <i>E. coli</i> , <i>Klebsiella</i> , <i>Proteus</i> , <i>Pseudomonas</i> sp. Enterococci	GNB GNB GPC
Intravenous catheter phlebitis and/or sepsis Peripheral catheter  Hyperalimentation line	<i>Staphylococcus aureus</i> or <i>S. epidermidis</i> <i>Klebsiella</i> , <i>Enterobacter</i> , <i>Pseudomonas</i> sp. <i>Candida</i> sp., <i>S. aureus</i> , <i>S. epidermidis</i> , enterococci <i>Klebsiella</i> , <i>Enterobacter</i> sp., etc.	GPC GNB Budding yeast; GPC GNB
Arteriovenous shunt	<i>S. aureus</i> , <i>S. epidermidis</i>	GPC
Septic bursitis	<i>S. aureus</i>	GPC
Biliary tract	<i>E. coli</i> , <i>Klebsiella</i> sp., and enterococci; <i>Bacteroides fragilis</i> (in elderly patients), <i>Clostridia</i> sp.	
Intra-abdominal abscess, peritonitis, or large bowel perforation; diverticulitis <sup>a</sup>	<i>E. coli</i> <i>B. fragilis</i>  <i>Klebsiella</i> sp. (Enterococci)	GNB GNB (thin, irregularly stained) GNB GPC
Burn wounds	Early: <i>S. aureus</i> , streptococci Later: Gram-negative bacilli, fungi	
Cellulitis, wound and soft tissue infections	<i>S. aureus</i> Streptococci	GPC GPC

Presumed location of Infection	Common pathogens	Gram stain Characteristics of exudate-if available
	<i>Clostridium</i> sp.	GPB
Meningitis	See Appendix C	
Pneumonia	See Appendix D	
Pelvic abscess, postabortal or postpartal	Anaerobic streptococci <i>B. fragilis</i>  <i>Clostridium</i> sp. <i>E. coli</i> Enterococci	GPC GNB (thin, irregularly stained) GPB GNB GPC
Septic arthritis	<i>S. aureus</i> <i>Haemophilus influenzae</i> (in children younger than 6 yr) Group B streptococci (in neonates) Gram-negative organisms <sup>b</sup>	GPC GNC  GPC GNB
Acute osteomyelitis	<i>S. aureus</i> <i>H. influenzae</i> (in children younger than 6 yr) Group B streptococci (in neonates) Gram-negative organisms <sup>b</sup>	GPC GNC  GPC GNB

Key: GNB = Gram-negative bacilli; GPC = Gram-positive cocci; GPB = Gram-positive bacilli; GNC = Gram-negative coccobacilli.

<sup>a</sup>The precise role of enterococci in intra-abdominal infections is unclear. In mild to moderate infections, it may not be necessary to provide antibiotic activity against enterococci.

<sup>b</sup>In high-risk patients (e.g., immunocompromised, elderly, IV drug abusers, diabetics, debilitated patients).

#### A. Dosage

To reduce the resistance of a microorganism to an antimicrobial agent, as exemplified by reducing the resistance of a bacterium to an antibiotic, or to kill a microorganism or bacterium, one would generally contact the microorganism or bacterium with an effective amount of the antibiotic or antimicrobial agent in

combination with an amount of an antimicrobial peptide effective to inhibit growth of the microorganism or bacterium. In terms of killing or reducing the resistance of a bacterium, one would contact the bacterium with an effective amount of an antibiotic in combination with an amount of an antimicrobial peptide effective to inhibit growth  
5 and/or proliferation in the bacterium. The terms "microorganism" and "bacterium" are used for simplicity and it will be understood that the invention is suitable for use against a population of microorganisms, *i.e.*, "bacteria".

The microorganism, *e.g.*, bacterium, or population thereof, may be contacted  
10 either *in vitro* or *in vivo*. Contacting *in vivo* may be achieved by administering to an animal (including a human patient) that has, or is suspected to have a microbial or bacterial infection, a therapeutically effective amount of pharmacologically acceptable antimicrobial peptide formulation in alone or in combination with a therapeutic amount of a pharmacologically acceptable formulation of an antibiotic agent. The invention may  
15 thus be employed to treat both systemic and localized microbial and bacterial infections by introducing the combination of agents into the general circulation or by applying the combination, *e.g.*, topically to a specific site, such as a wound or burn, or to the eye, ear or other site of infection.

20 Where an antimicrobial peptide is used in combination with other antimicrobial agents or antibiotics, an "effective amount of an antimicrobial agent or antibiotic" means an amount, or dose, within the range normally given or prescribed. Such ranges are well established in routine clinical practice and will thus be known to those of skill in the art. Appropriate oral and parenteral doses and treatment regimens are further detailed herein  
25 in Table 5 and Table 6. As this invention provides for enhanced microbial and/or bacterial killing, it will be appreciated that effective amounts of an antimicrobial agent or antibiotic may be used that are lower than the standard doses previously recommended when the antimicrobial or antibiotic is combined with a antimicrobial peptide.

30 Naturally, in confirming the optimal therapeutic dose for antimicrobial peptides, first animal studies and then clinical trials would be conducted, as is routinely practiced

in the art. Animal studies are common in the art and are further described herein (Example II) and in publications such as Lorian (1991, pp. 746-786, incorporated herein by reference) and Cleeland and Squires (incorporated herein by reference, from within the Lorian text).

5

The ID<sub>50</sub>/IC<sub>50</sub> ratio required for safe use of the proposed inhibitor-antimicrobial peptide or combinations of peptide with other antimicrobial agents will be assessed by determining the ID<sub>50</sub> (median lethal toxic dosage) and the IC<sub>50</sub> (median effective therapeutic dosage) in experimental animals. The optimal dose for human subjects is then defined by fine-tuning the range in clinical trials. In the case of ID<sub>50</sub>, the inhibitor is usually administered to mice or rats (orally or intraperitoneal) at several doses (usually 4-5) in the lethal range. The dose in mg/kg is plotted against % mortality and the dose at 50% represents the ID<sub>50</sub> (Klaassen, 1990). The IC<sub>50</sub> is determined in a similar fashion as described by Cleeland and Squires (1991).

15

In a clinical trial, the therapeutic dose would be determined by maximizing the benefit to the patient, whilst minimizing any side-effects or associated toxicities. Throughout the detailed examples, various therapeutic ranges are listed. Unless otherwise stated, these ranges refer to the amount of an agent to be administered orally.

20

In optimizing a therapeutic dose within the ranges disclosed herein, one would not use the upper limit of the range as the starting point in a clinical trial due to patient heterogeneity. Starting with a lower or mid-range dose level, and then increasing the dose will limit the possibility of eliciting a toxic or untoward reaction in any given patient or subset of patients. The presence of some side-effects or certain toxic reactions *per se* would not, of course, limit the utility of the invention, as it is well known that most beneficial drugs also produce a limited amount of undesirable effects in certain patients. Also, a variety of means are available to the skilled practitioner to counteract certain side-effects, such as using vitamin B<sub>12</sub> in association with N<sub>2</sub>O treatment (Ostreicher, 1994).

30

Zak and Sande (1981) reported on the correlation between the *in vitro* and *in vivo* activity of a 1000 compounds that were randomly screened for antimicrobial activity. The important finding in this study is that negative *in vitro* data is particularly accurate, with the negative *in vitro* results showing more than a 99% correlation with negative  
5 *in vivo* activity. This is meaningful in the context of the present invention as one or more *in vitro* assays will be conducted prior to using any given combination in a clinical setting. Any negative result obtained in such an assay will thus be of value, allowing efforts to be more usefully directed.

10 In the treatment of animals or human patients with combination therapy, there are various appropriate formulations and treatment regimens that may be used. For example, the antimicrobial peptide and second agent(s) may be administered to an animal simultaneously, *e.g.*, in the form of a single composition that includes the antimicrobial peptide and second agent, or by using at least two distinct compositions. The  
15 antimicrobial agent could also be administered to the animal prior to the second agent or the second agent may be given prior to the antimicrobial peptide.

Multiple combinations may also be used, such as more than one antimicrobial peptide used with one second agent or more than one second agent. Different classes  
20 second agents and antimicrobial peptides may be combined, naturally following the general guidelines known in the art regarding drug interactions. Typically, between one and about five distinct antimicrobial agents are contemplated for use along with between one and about six antimicrobial peptides.

25 Further embodiments of the invention include therapeutic kits that comprise, in suitable container means, a pharmaceutical formulation of at least one antimicrobial peptide and a pharmaceutical formulation of at least one antimicrobial agent or antibiotic. The antimicrobial peptide and antimicrobial agent or antibiotic may be contained within a single container means, or a plurality of distinct containers may be employed.

30



Depending on the circumstances, antimicrobial agents may be employed in oral or parenteral treatment regimens. Appropriate doses are well known to those of skill in the art and are described in various publications, such as (Reese and Betts, 1993; incorporated herein by reference). Table 5 and Table 6 (taken from Reese and Betts, 1993) are included herein to provide ready reference to the currently recommended doses of a variety of antimicrobial agents.

Following are definitions of terms that are used in Table 5 and Table 6: qid (4 times daily), tid (3 times daily), bid (twice daily), qd (once daily), q4h (every 4 hours around the clock), q6h (every 6 hours around the clock) and q8h (every 8 hours around the clock).

**TABLE 5:**  
**COMMON ANTIBIOTICS AND USUAL ORAL DOSES**

ANTIBIOTIC	DOSAGE
Penicillin V Rugby (generic) V-cillin K	50 mg qid <sup>2</sup>
Dicloxacillin Glenlawn (generic) Dynapen	250 mg qid
Cloxacillin (Tegopen)	250 mg qid
Amoxicillin Rugby (generic) Polymox	250 mg tid
Ampicillin Moore (generic) Polycillin	250 mg qid
Augmentin 250-mg tablets chewables (250 mg) 125-mg (suspension) chewables (125 mg)	tid
Carbenicillin (Geocillin)	382 mg qid (1 tb) 2 tab qid

ANTIBIOTIC	DOSAGE
Cephalexin Rugby (generic) Keflex	250 mg qid
Rugby (generic) Keflex	500 mg qid
Cefadroxil Rugby (generic) Duricef	1 gm bid
Cephradine Rugby (generic) Velosef	250 mg qid
Rugby (generic) Velosef	500 mg qid
Cefaclor Ceclor	250 mg tid
Cefuroxime axetil Ceftin	125 mg bid 250 mg bid 500 mg bid
Cefixime Suprax	400 mg q24h
Cefprozil Cefzil	250 mg q12h
Loracarbef (Lorabid)	200 mg bid
Cefpodoxime proxetil (Vantin)	200 mg bid
Clindamycin Cleocin	300 mg q8h
TMP/SMZ Bactrim Septra (generic)	1 double-strength bid
Trimethoprim Rugby (generic) Proloprim	100 mg bid
Erythromycin (base) Abbott	250 mg qid

ANTIBIOTIC	DOSAGE
E-mycin (delayed release)	
Erythromycin stearate Rugby (generic)	250 mg qid
Azithromycin Zithromax	1 g once only 500 mg, day 1, plus 250 mg, day 2-5
Clarithromycin Biaxin	250 mg bid 500 mg bid
Tetracycline hydrochloride Mylan Sumycin 250	250 mg qid
Doxycycline  Lederle (generic) Vibramycin	100 mg qd (with 200- mg initial load)
Vancomycin Vancocin HCl (oral soln/powder)	Capsules 125 mg q6h PO
Metronidazole Rugby (generic) Flagyl	250 mg qid
Norfloxacin Noroxin	400 mg bid
Ciprofloxacin Cipro	250 mg bid 500 mg bid 750 mg bid
Ofloxacin Floxin	200 mg bid 300 mg bid 400 mg bid
Lomefloxacin Maxaquin	400 mg once qd

**TABLE 6:**  
**COMMON ANTIBIOTICS AND USUAL PARENTERAL DOSES**

ANTIBIOTIC	DOSAGE
Penicillin G Pfizerpen G (Pfizer)	2,400,000 units 12 million units
Oxacillin Prostaphlin (Bristol)	12 g
Nafcillin Nafcil (Bristol)	12 g
Ampicillin Omnipen (Wyeth)	6 g
Ticarcillin Ticar (Beecham)	18 g
Piperacillin Pipracil (Lederle)	18 g 16 g
Mezlocillin Mezlin (Miles)	18 g 16 g
Ticarcillin-clavulanate Timentin (Beecham)	18 g/0.6 g 12 g/0.4 g
Ampicillin-sulbactam Unasyn (Roerig)	6 g 12 g
Cephalothin Keflin (Lilly)	9 g (1.5 g q4h)
Cefazolin Ancef (SKF)	4 g (1 g q6h) 3 g (1 g q8h)
Cefuroxime Zinacef (Glaxo)	6 g 2.25 g (750 mg q8h) 4.5 g (1.5 g q8h)
Cefamandole Mandol (Lilly)	9 g (1.5 g q4h)
Cefoxitin Mefoxin (MSD)	8 g (2 g q6h) 6 g (2 g q8h)
Cefonicid Monicid (SKF)	1 g q12h
Cefotetan Cefotan (Stuart)	2 g q12h
Cefmetazole Zefazone (Upjohn)	2 g q8h

ANTIBIOTIC	DOSAGE
Ceftriaxone Rocephin (Roche)	2 g (2.0 g q24h) 1 g (1.0 g q24h)
Ceftazidime Fortax (Glaxo) Taxicef (SKF) Tozidime (Lilly)	6 g (2 g q8h)
Cefotaxime Claforan (Hoechst)	2 g q6h 2 g q8h
Cefoperazone Cefobid (Pfizer)	8 g (2 g q6h) 6 g (2 g q8h)
Ceftizoxime Ceftizox (SKF)	(2 g q8h)
Aztreonam Azactam (Squibb)	2 g q8h 1 g q8h
Imipenem Primaxin (MSD)	2000 mg (500 mg 16h)
Gentamicin Garamycin (Schering) (generic) (Elkins-Sinn)	360 mg (1.5 mg/kg q8h for an 80-kg patient)
Tobramycin Nebcin (Dista)	360 mg (1.5 mg/kg q8h for an 80-kg patient)
Amikacin Amikin (Bristol)	1200 mg (7.5 mg/kg q12h for an 80-kg patient)
Clindamycin Cleocin (Upjohn)	2400 mg (600 mg q6h) 2700 mg (900 mg q8h) 1800 mg (600 mg q8h)
Chloramphenicol Chloromycetin (P/D)	4 g (1 g q6h)
TMP/SMZ Septra (Burroughs Wellcom)	1400 mg TMP (5 mg TMP/kg q6h for a 70-kg patient) 700 mg TMP (5 mg TMP/kg q12h for a 70- kg patient)
Erythromycin	2000 mg (500 mg q6h)

ANTIBIOTIC	DOSAGE
Erythromycin (Elkins-Sinn)	
Doxycycline Vibramycin (Pfizer)	200 mg (100 mg q12h)
Vancomycin Vancocin (Lilly)	2000 mg (500 mg q6h)
Metronidazole (generic) (Elkins-Sinn)	2000 mg (500 mg q6h)
Ciprofloxacin Cipro	200 mg q12h 400 mg q12h
Pentamidine Pentam (LyphoMed)	280 mg (4 mg/kg q24h for a 70-kg patient)

5 The effectiveness of erythromycin and lincomycin against a wide variety of organisms is shown in Table 7 (taken from Lorian, 1991) to illustrate the range of antibiotic resistance acquired by various bacterial strains. The data presented in the tables of the present specification is merely illustrative and is considered another tool to enable the straightforward comparison of raw data with accepted clinical practice and to allow the determination of appropriate doses of combined agents for clinical use.

10

TABLE 7:  
SUSCEPTIBILITY TO ANTIBIOTICS

ERYTHROMYCIN

Species	(n)	e	Rang	MIC <sub>50</sub>	MIC <sub>90</sub>
<i>Bacillus</i> spp.	20	0.03-2		0.25	2
<i>Bacteroides fragilis</i>	97	0.25-16		1	8
<i>Bordetella bronchiseptica</i>	11	4-32		8	32
<i>Bordetella parapertussis</i>	46	0.125-4		0.25	0.25
<i>Bordetella pertussis</i>	32	1-0.5		0.25	0.25
<i>Bordetella pertussis</i>	75	0.125-0.5		0.125	0.125
<i>Borrelia burgdorferi</i>	10	0.03-0.125		0.03	0.06



Species		(n)	Rang e	MIC <sub>50</sub>	MIC <sub>90</sub>
<i>Branhamella</i> <i>catarrhalis</i>	(Moraxella)	20	0.125-0.5	0.25	0.25
<i>Branhamella</i> <i>catarrhalis</i>	(Moraxella)	20	0.125-0.5	0.25	1
<i>Branhamella</i> <i>catarrhalis</i> (non producer)	(Moraxella) $\beta$ -lactamase	40	0.06-0.5	0.25	0.5
<i>Branhamella</i> <i>catarrhalis</i> (non $\beta$ -lactamase producer)	(Moraxella)	13	0.03-0.125	0.06	0.06
<i>Branhamella</i> <i>catarrhalis</i> (non producer)	(Moraxella) $\beta$ -lactamase	14	0.06-1	0.125	1
<i>Branhamella</i> <i>catarrhalis</i> (non producer)	(Moraxella) $\beta$ -lactamase	16	0.015-1	0.06	0.25
<i>Branhamella</i> <i>catarrhalis</i> ( $\beta$ -lactamase producer)	(Moraxella)	47	0.06-1	0.25	0.5
<i>Branhamella</i> <i>catarrhalis</i> ( $\beta$ -lactamase producer)	(Moraxella)	58	0.03-0.25	0.125	0.125
<i>Branhamella</i> <i>catarrhalis</i> ( $\beta$ -lactamase producer)	(Moraxella)	160	0.06-8	0.25	0.5
<i>Branhamella</i> <i>catarrhalis</i> ( $\beta$ -lactamase producer)	(Moraxella)	35	0.03-0.125	0.06	0.06
<i>Campylobacter jejuni</i>		25	0.5-8	1	4
<i>Campylobacter jejuni</i>		16	0.125-4	0.25	2
<i>Campylobacter pylori</i>		56	0.25-16	0.5	1
<i>Campylobacter pylori</i>		13	0.125-0.25	0.125	0.25
<i>Corynebacterium JK</i>		102	0.5- <sup>3</sup> 128	<sup>3</sup> 128	<sup>3</sup> 128
<i>Corynebacterium JK</i>		19	0.125- <sup>3</sup> 64	2	<sup>3</sup> 64
<i>Enterococcus faecalis</i>		26	1- <sup>3</sup> 64	1	4
<i>Enterococcus faecalis</i>		50	0.06- <sup>3</sup> 64	4	<sup>3</sup> 64

Species	(n)	Rang e	MIC <sub>50</sub>	MIC <sub>90</sub>
<i>Enterococcus faecalis</i>	86	0.125- <sup>3</sup> 64	1	<sup>3</sup> 64
<i>Enterococcus faecalis</i>	97	0.125-128	2	128
<i>Enterococcus faecium</i>	14	0.06- <sup>3</sup> 64	1	<sup>3</sup> 64
<i>Enterococcus</i> spp.	35	0.06- <sup>3</sup> 32	2	<sup>3</sup> 32
<i>Haemophilus ducreyi</i>	122	?-0.125	0.004	0.06
<i>Haemophilus influenzae</i>	145	0.5-8	2	2
<i>Haemophilus influenzae</i>	97	0.25- <sup>3</sup> 16	1	4
<i>Haemophilus influenzae</i> (non $\beta$ -lactamase producer)	22	0.125-8	2	4
<i>Haemophilus influenzae</i> (non $\beta$ -lactamase producer)	137	0.06- <sup>3</sup> 8	4	8
<i>Haemophilus influenzae</i> ( $\beta$ -lactamase producer)	46	0.06-8	4	8
<i>Haemophilus influenzae</i> ( $\beta$ -lactamase producer)	17	0.25-4	2	4
<i>Haemophilus influenzae</i> (penicillin susceptible)	22	0.25-16	8	16
<i>Haemophilus influenzae</i> (penicillin resistant)	20	8-16	8	16
<i>Haemophilus parainfluenzae</i>	13	0.5-8	2	4
<i>Legionella</i> spp.	23	0.03-0.25	0.125	0.25
<i>Legionella pneumophila</i>	31	0.0075-0.25	0.06	0.125
<i>Legionella pneumophila</i>	48	0.03-2	0.25	0.5
<i>Legionella pneumophila</i>	25	0.125-1	0.25	1
<i>Listeria monocytogenes</i>	13	0.5-1	0.5	0.5
<i>Listeria monocytogenes</i>	16	0.125-2	0.25	1
<i>Listeria monocytogenes</i>	65	0.06- <sup>3</sup> 32	0.125	32
<i>Mycoplasma hominis</i>	26	<sup>3</sup> 128	<sup>3</sup> 128	<sup>3</sup> 128
<i>Mycoplasma hominis</i>	20	<sup>3</sup> 256	<sup>3</sup> 256	<sup>3</sup> 256
<i>Mycoplasma pneumoniae</i>	10	0.06-8	0.06	0.06

Species	(n)	Rang e	MIC <sub>50</sub>	MIC <sub>90</sub>
<i>Mycoplasma pneumoniae</i>	14	0.004-0.03	0.004	0.004
<i>Neisseria gonorrhoeae</i>	19	0.0075-8	0.25	1
<i>Neisseria gonorrhoeae</i> (non $\beta$ -lactamase producer)	73	0.015-4	0.25	2
<i>Neisseria gonorrhoeae</i> (non $\beta$ -lactamase producer)	78	0.03-2	0.25	1
<i>Neisseria gonorrhoeae</i> ( $\beta$ -lactamase producer)	12	0.03-4	0.5	2
<i>Neisseria gonorrhoeae</i> ( $\beta$ -lactamase producer)	17	1-4	2	4
<i>Neisseria meningitidis</i>	19	0.5-8	1	8
<i>Nocardia asteroides</i>	78	0.25- <sup>3</sup> 8	<sup>3</sup> 8	<sup>3</sup> 8
<i>Staphylococcus aureus</i>	44	0.125-1	0.125	0.5
<i>Staphylococcus aureus</i>	100	0.25-128	0.5	4
<i>Staphylococcus aureus</i> (penicillin susceptible)	20	0.125-0.5	0.5	0.5
<i>Staphylococcus aureus</i> (penicillin susceptible)	35	0.06- <sup>3</sup> 32	0.25	0.5
<i>Staphylococcus aureus</i> (penicillin resistant)	35	0.25- <sup>3</sup> 32	0.25	<sup>3</sup> 32
<i>Staphylococcus aureus</i> (methicillin susceptible)	28	0.125-1	0.25	0.5
<i>Staphylococcus aureus</i> (methicillin susceptible)	97	0.125- <sup>3</sup> 64	0.25	<sup>3</sup> 64
<i>Staphylococcus aureus</i> (methicillin susceptible)	20	0.125-1	0.5	0.5
<i>Staphylococcus aureus</i> (methicillin resistant)	17	0.5- <sup>3</sup> 128	128	128
<i>Staphylococcus aureus</i> (methicillin resistant)	15	<sup>3</sup> 64	<sup>3</sup> 64	<sup>3</sup> 64
<i>Staphylococcus aureus</i> (methicillin resistant)	20	<sup>3</sup> 64	<sup>3</sup> 64	<sup>3</sup> 64

Species	(n)	Rang e	MIC <sub>50</sub>	MIC <sub>90</sub>
<i>Staphylococcus aureus</i> (methicillin resistant)	30	0.06- <sup>3</sup> 32	<sup>3</sup> 32	<sup>3</sup> 32
<i>Staphylococcus coagulase f</i>	10	0.125-4	0.25	2
<i>Staphylococcus coagulase f</i>	100	0.125- <sup>3</sup> 64	0.25	<sup>3</sup> 64
<i>Staphylococcus coagulase f</i> (non $\beta$ -lactamase producer)	12	0.03-8	0.125	0.25
<i>Staphylococcus coagulase f</i> ( $\beta$ -lactamase producer)	38	0.06-16	0.125	4
<i>Staphylococcus epidermidis</i>	50	0.125- <sup>3</sup> 64	<sup>3</sup> 64	<sup>3</sup> 64
<i>Staphylococcus haemolyticus</i>	20	0.125- <sup>3</sup> 64	<sup>3</sup> 64	<sup>3</sup> 64
<i>Staphylococcus hominis</i>	20	0.125- <sup>3</sup> 64	<sup>3</sup> 64	<sup>3</sup> 64
<i>Streptococcus agalactiae</i>	20	0.03-0.25	0.03	0.125
<i>Streptococcus agalactiae</i>	34	0.015-0.06	0.03	0.03
<i>Streptococcus pneumoniae</i>	58	0.03-0.25	0.06	0.125
<i>Streptococcus pneumoniae</i>	91	0.125-4	0.125	0.125
<i>Streptococcus pneumoniae</i>	50	0.015-0.06	0.03	0.03
<i>Streptococcus pneumoniae</i>	16	0.03-0.125	0.06	0.125
<i>Streptococcus pneumoniae</i>	26	0.015-0.25	0.03	0.06
<i>Streptococcus pneumoniae</i>	50	0.03-0.125	0.06	0.06
<i>Streptococcus pyogenes</i>	19	0.03-0.25	0.06	0.125
<i>Streptococcus pyogenes</i>	20	0.03-0.25	0.06	0.125
<i>Streptococcus pyogenes</i>	33	0.015-0.03	0.03	0.03
<i>Streptococcus pyogenes</i>	20	0.06- <sup>3</sup> 32	0.125	<sup>3</sup> 32
<i>Streptococcus spp.</i>	22	0.015-0.25	0.03	0.06
<i>Streptococcus spp.</i>	107	0.004-2	0.03	1
<i>Ureaplasma urealyticum</i>	28	0.015- <sup>3</sup> 256	2	<sup>3</sup> 256
<i>Ureaplasma urealyticum</i>	19	8- <sup>3</sup> 128	16	32

# LINCOMYCIN

Species	(n)	Rang	MIC <sub>50</sub>	MIC <sub>90</sub>
<i>Mycoplasma hominis</i>	28	0.5-16	2	4
<i>Mycoplasma pneumoniae</i>	11	2-32	8	32
<i>Staphylococcus aureus</i>	100	0.5-512	1	1
<i>Ureaplasma urealyticum</i>	19	64- <sup>3</sup> 128	<sup>3</sup> 128	<sup>3</sup> 128

#### B. *Pharmaceutical Compositions*

The phrases "pharmaceutically" or "pharmacologically acceptable" refer to molecular entities and compositions that do not produce adverse, allergic, or other untoward reactions when administered to an animal or a human. As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the vectors or cells of the present invention, its use in therapeutic compositions is contemplated. Supplementary active ingredients also can be incorporated into the compositions.

The active compositions of the present invention may include classic pharmaceutical preparations. Administration of these compositions according to the present invention will be via any common route so long as the target tissue is available via that route. This includes oral, nasal, buccal, rectal, vaginal or topical. Alternatively, administration may be by orthotopic, intradermal, subcutaneous, intramuscular, intraperitoneal or intravenous injection. Such compositions would normally be administered as pharmaceutically acceptable compositions, described *supra*.

The active compounds may also be administered parenterally or intraperitoneally. Solutions of the active compounds as free base or pharmacologically acceptable salts can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose.

Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

5           The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of  
10 microorganisms, such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of  
15 dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the  
20 compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

          Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients  
25 enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-  
30 drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.



As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

For oral administration the polypeptides of the present invention may be incorporated with excipients and used in the form of non-ingestible mouthwashes and dentifrices. A mouthwash may be prepared incorporating the active ingredient in the required amount in an appropriate solvent, such as a sodium borate solution (Dobell's Solution). Alternatively, the active ingredient may be incorporated into an antiseptic wash containing sodium borate, glycerin and potassium bicarbonate. The active ingredient may also be dispersed in dentifrices, including: gels, pastes, powders and slurries. The active ingredient may be added in a therapeutically effective amount to a paste dentifrice that may include water, binders, abrasives, flavoring agents, foaming agents, and humectants.

The compositions of the present invention may be formulated in a neutral or salt form. Pharmaceutically-acceptable salts include the acid addition salts (formed with the free amino groups of the protein) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like.

Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective. The formulations

are easily administered in a variety of dosage forms such as injectable solutions, drug release capsules and the like. Routes of administration may be selected from intravenous, intrarterial, intrabuccal, intraperitoneal, intramuscular, subcutaneous, oral, topical, rectal, vaginal, nasal and intraocular.

5

For parenteral administration in an aqueous solution, for example, the solution should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, sterile aqueous media which can be employed will be known to those of skill in the art in light of the present disclosure. For example, one dosage could be dissolved in 1 ml of isotonic NaCl solution and either added to 1000 ml of hypodermoclysis fluid or injected at the proposed site of infusion, (see for example, "Remington's Pharmaceutical Sciences" 15th Edition, pages 1035-1038 and 1570-1580). Some variation in dosage will necessarily occur depending on the condition of the subject being treated. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject. Moreover, for human administration, preparations should meet sterility, pyrogenicity, general safety and purity standards as required by FDA Office of Biologics standards.

20

In a particular embodiment, liposomal formulations are contemplated. Liposomal encapsulation of pharmaceutical agents prolongs their half-lives when compared to conventional drug delivery systems. Because larger quantities can be protectively packaged, this allow the opportunity for dose-intensity of agents so delivered to cells. This would be particularly attractive in the chemotherapy of cervical cancer if there were mechanisms to specifically enhance the cellular targeting of such liposomes to these cells.

25

"Liposome" is a generic term encompassing a variety of single and multilamellar lipid vehicles formed by the generation of enclosed lipid bilayers. Phospholipids are used for preparing the liposomes according to the present invention and can carry a net positive

30

charge, a net negative charge or are neutral. Dicetyl phosphate can be employed to confer a negative charge on the liposomes, and stearylamine can be used to confer a positive charge on the liposomes. Liposomes are characterized by a phospholipid bilayer membrane and an inner aqueous medium. Multilamellar liposomes have multiple lipid layers separated by aqueous medium. They form spontaneously when phospholipids are suspended in an excess of aqueous solution. The lipid components undergo self-rearrangement before the formation of closed structures and entrap water and dissolved solutes between the lipid bilayers (Ghosh and Bachhawat, 1991). Also contemplated are cationic lipid-nucleic acid complexes, such as lipofectamine-nucleic acid complexes.

10

In certain embodiments of the invention, the liposome may be complexed with a hemagglutinating virus (HVJ). This has been shown to facilitate fusion with the cell membrane and promote cell entry of liposome-encapsulated DNA (Kaneda *et al.*, 1989). In other embodiments, the liposome may be complexed or employed in conjunction with nuclear non-histone chromosomal proteins (HMG-1) (Kato *et al.*, 1991). In yet further embodiments, the liposome may be complexed or employed in conjunction with both HVJ and HMG-1. In that such expression vectors have been successfully employed in transfer and expression of a polynucleotide *in vitro* and *in vivo*, then they are applicable for the present invention. Where a bacterial promoter is employed in the DNA construct, it also will be desirable to include within the liposome an appropriate bacterial polymerase.

20

Lipids suitable for use according to the present invention can be obtained from commercial sources. For example, dimyristyl phosphatidylcholine ("DMPC") can be obtained from Sigma Chemical Co., dicetyl phosphate ("DCP") is obtained from K & K Laboratories (Plainview, NY); cholesterol ("Chol") is obtained from Calbiochem-Behring; dimyristyl phosphatidylglycerol ("DMPG") and other lipids may be obtained from Avanti Polar Lipids, Inc. (Birmingham, Ala.). Stock solutions of lipids in chloroform, chloroform/methanol or *t*-butanol can be stored at about -20°C. Preferably, chloroform is used as the only solvent since it is more readily evaporated than methanol.

30

Phospholipids from natural sources, such as egg or soybean phosphatidylcholine, brain phosphatidic acid, brain or plant phosphatidylinositol, heart cardiolipin and plant or bacterial phosphatidylethanolamine are preferably not used as the primary phosphatide, *i.e.*, constituting 50% or more of the total phosphatide composition, because of the instability and leakiness of the resulting liposomes.

Liposomes used according to the present invention can be made by different methods. The size of the liposomes varies depending on the method of synthesis. A liposome suspended in an aqueous solution is generally in the shape of a spherical vesicle, having one or more concentric layers of lipid bilayer molecules. Each layer consists of a parallel array of molecules represented by the formula XY, wherein X is a hydrophilic moiety and Y is a hydrophobic moiety. In aqueous suspension, the concentric layers are arranged such that the hydrophilic moieties tend to remain in contact with an aqueous phase and the hydrophobic regions tend to self-associate. For example, when aqueous phases are present both within and without the liposome, the lipid molecules will form a bilayer, known as a lamella, of the arrangement XY-YX.

Liposomes within the scope of the present invention can be prepared in accordance with known laboratory techniques. In one preferred embodiment, liposomes are prepared by mixing liposomal lipids, in a solvent in a container, *e.g.*, a glass, pear-shaped flask. The container should have a volume ten-times greater than the volume of the expected suspension of liposomes. Using a rotary evaporator, the solvent is removed at approximately 40°C under negative pressure. The solvent normally is removed within about 5 min to 2 hours, depending on the desired volume of the liposomes. The composition can be dried further in a desiccator under vacuum. The dried lipids generally are discarded after about 1 week because of a tendency to deteriorate with time.

Dried lipids can be hydrated at approximately 25-50 mM phospholipid in sterile, pyrogen-free water by shaking until all the lipid film is resuspended. The aqueous liposomes can be then separated into aliquots, each placed in a vial, lyophilized and sealed under vacuum.

In the alternative, liposomes can be prepared in accordance with other known laboratory procedures: the method of Bangham *et al.* (1965), the contents of which are incorporated herein by reference; the method of Gregoriadis (1979), the contents of which are incorporated herein by reference; the method of Deamer and Uster (1983), the contents of which are incorporated by reference; and the reverse-phase evaporation method as described by Szoka and Papahadjopoulos (1978). The aforementioned methods differ in their respective abilities to entrap aqueous material and their respective aqueous space-to-lipid ratios.

10

The dried lipids or lyophilized liposomes prepared as described above may be reconstituted in a solution of nucleic acid and diluted to an appropriate concentration with an suitable solvent, *e.g.*, DPBS. The mixture is then vigorously shaken in a vortex mixer. Unencapsulated nucleic acid is removed by centrifugation at  $29,000 \times g$  and the liposomal pellets washed. The washed liposomes are resuspended at an appropriate total phospholipid concentration, *e.g.*, about 50-200 mM. The amount of nucleic acid encapsulated can be determined in accordance with standard methods. After determination of the amount of nucleic acid encapsulated in the liposome preparation, the liposomes may be diluted to appropriate concentration and stored at 4°C until use.

20

In a preferred embodiment, the lipid dioleoylphosphatidylchoine is employed. Nuclease-resistant oligonucleotides were mixed with lipids in the presence of excess *t*-butanol. The mixture was vortexed before being frozen in an acetone/dry ice bath. The frozen mixture was lyophilized and hydrated with Hepes-buffered saline (1 mM Hepes, 10 mM NaCl, pH 7.5) overnight, and then the liposomes were sonicated in a bath type sonicator for 10 to 15 min. The size of the liposomal-oligonucleotides typically ranged between 200-300 nm in diameter as determined by the submicron particle sizer autodilute model 370 (Nicomp, Santa Barbara, CA).

30

The purified antimicrobial peptide may be used without further modifications or it may be diluted in a pharmaceutically acceptable carrier. The peptides may be used



independently or in combination with other antimicrobial agents. Because of the stability of the peptides it is contemplated that the invention may be administered to humans or animals, included in food preparations, pharmaceutical preparations, medicinal and pharmaceutical products, cosmetic products, hygienic products, cleaning products and  
5 cleaning agents, as well as any material to which the peptides could be sprayed on or adhered to wherein the inhibition of microbial growth is desired.

### 3. Examples

The following examples are included to demonstrate preferred embodiments of  
10 the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific  
15 embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

#### EXAMPLE 1: Peptide Synthesis

All peptides were synthesized by the solid-phase method employing an Applied  
20 Biosystems model 433A peptide synthesizer and Fastmoc strategy at the 0.1 mM scale. Peptides were purified by reversed-phase HPLC on a Waters Delta Prep employing a Vydac 218TP1022 (22 x 250 mm) column. Separation was performed with a gradient system of aqueous 0.1% trifluoroacetic acid (solvent A) and 100% acetonitrile containing 0.085% trifluoroacetic acid (solvent B). A linear gradient from 0 to 100% B was applied  
25 over 70 min and fractions collected every 0.2 min. Fractions were subsequently monitored by analytical scale reversed-phase HPLC on a Beckman Gold System using a Vydac 218TP54 (4.6 x 250 mm) column at a flow rate of 0.5 ml/min under isocratic elution conditions. Select fractions were pooled and lyophilized; further characterization of peptides was provided by mass spectrometry and capillary electrophoresis. Mass  
30 measurements were performed by flow injection at 0.1 ml/min in 64% acetonitrile containing 0.05% trifluoroacetic acid with a Hewlett-Packard model 1100 MSD equipped



with an electrospray ionization source. Capillary electrophoresis was performed on a Hewlett-Packard 3D instrument equipped with an extended light path fused-silicate column 75 micrometers (ID) x 80.5 centimeters (total length). Capillary electrophoresis experiments were conducted at 18 C in 100 mM sodium phosphate buffer, pH 2.9 at  
5 20,000 volts. Peptide concentration was determined by quantitative amino acid analysis on a Beckman 6300 Amino Acid analyzer.

### Example 2: Bacterial Strains and Antimicrobial Assays

We adapted a luminescence assay that we had previously used with *E. coli* (Singh,  
10 1999 and Travis, 1999) to examine the time-course of killing and the potency of the peptides against *P. aeruginosa* PAO1 in high and low salt buffer solutions. This assay measured antimicrobial activity as a decrease in energy-dependent luminescence of bacteria containing luminescence genes. *P. aeruginosa* strain PAO1 was transformed with the *Vibrio fischeri* luminescence genes on the pBBR1MCS-5 plasmid (Frackman,  
15 1999). *P. aeruginosa* was grown to early log phase ( $2.5 \times 10^7$  bacteria/ml) at 30°C in tryptic soy broth containing 25 µg/ml gentamicin, then used without centrifugation. For each assay,  $5 \times 10^5$  bacteria were incubated in 6.7 mM potassium phosphate, pH 7.4, 3.3% tryptic soy broth, antimicrobial peptide and other additions in a total volume of 150 µl in 96-well plates (Optiplate, Packard Instruments). After incubation at 30°C for the  
20 indicated time, relative emission of light (in arbitrary units) was measured with a Packard Instruments luminometer (Meriden, Connecticut). We defined the EC<sub>50</sub> as the amount of antimicrobial peptide that decreased luminescence by 50%. The ionic strength of the standard low salt assay solution was calculated to be 25 mM. In some experiments the ionic strength was increased to 175 mM by the addition of 150 mM NaCl. For colony  
25 counting assays, bacteria were prepared and incubated with antimicrobial peptides as described above, then plated on nutrient agar plates. Colonies were counted after incubation of the plates for 24-48 hr at 37°C.

Minimal inhibitory concentrations (MICs) for Gram-positive and Gram-negative  
30 clinical isolates were determined by a Microbroth tray dilution assay. Serial two-fold dilutions of peptides (0.06-32 µg/ml) in Mueller-Hinton broth ([Na<sup>+</sup>] = 128 mM, [Cl<sup>-</sup>] =

108 mM) were inoculated with 100,000 CFU of bacteria, incubated at 35°C and read at 24 hr.

The two stage radial diffusion assay used in these studies has been described elsewhere (Lehrer et al., 1991). Briefly, the purified peptides were serially diluted in acidified water (0.01% acetic acid) that contained 0.1 % human serum albumin (Sigma A-8763). The bacteria (*E. coli* ML-35p, *P. aeruginosa* MR 3007, *S. aureus* 930918-3, and methicillin-resistant *S. aureus* ATCC 33591 (MRSA) were grown to mid-logarithmic phase in trypticase soy broth and washed with 10 mM phosphate buffer, pH 7.4. Approximately  $2 \times 10^5$  CFU/ml was incorporated into a thin (1.23 mm) agarose underlay gel that contained 1% w/v agarose (Sigma A-6013), 10 mM sodium phosphate buffer, pH 7.4, and 0.3 mg/ml trypticase soy broth powder  $\pm$  100 mM NaCl. A regularly spaced, 5-by-5 array of wells was made in the underlay gel. The wells, 3.2 mm in diameter, had a 10  $\mu$ l capacity. Six serially diluted samples of each peptide were prepared, ranging in concentration from 0.79 to 250  $\mu$ g/ml, and 5  $\mu$ l aliquots were added to the wells. After 3 hours, a 10 ml overlay gel composed of 6% trypticase soy broth powder, 1% agarose and 10 mM sodium phosphate buffer, pH 7.4 was poured, and the plates were incubated overnight to allow surviving organisms to form microcolonies.

The zone diameters were measured to the nearest 0.1 mm and were expressed in units (1 unit = 0.1 mm), after first subtracting the diameter of the well. A linear relationship existed between the zone diameter and the  $\log_{10}$  of the peptide concentration. The minimal inhibitory concentration was determined by performing a least mean squares fit and solving for the X-intercept with a Hewlett Packard 20S Scientific Calculator or its equivalent.

### Example 3: CD Spectroscopy

Circular dichroism (CD) spectroscopy was performed on an AVIV 62 DS spectrometer (AVIV Associates, Lakewood, NJ) equipped with a thermoelectric temperature control at 25 C. Samples contained 0.11 - 0.24 mg/mL peptide in either 50 mM sodium phosphate, pH 7.0; some samples also contained 40% trifluoroethanol or

0.1% (0.22 mM) lipopolysaccharide (LPS). LPS from a clinical isolate of *Neisseria meningitidis* Type B was provided by Dr. M. A. Apicella. Spectra were collected at 0.5 nm intervals with an averaging time of 2 s per datum point using a pathlength of 0.1 cm. The spectra were smoothed once over an interval of five data points prior to plotting with  
5 each spectrum representing the average of two scans. The average of two buffer scans was subtracted prior to data smoothing. The fractional helical content was calculated from the ratio of observed mean residue ellipticity to the mean residue ellipticity for a 100% helical peptide of identical length at 25°C according to Luo and Baldwin (1997).

#### 10 **Example 4: Calculation of Peptide Hydrophobic Moment and Hydrophobicity**

Peptide hydrophobicities and hydrophobic moments were calculated as described by Eisenberg (1986) using normalized consensus hydrophobicity scales.

#### **Example 5: Hemolysis Assay**

15 Hemolytic activity of the peptides was assayed with heparinized human red blood cells. The cells were collected from a normal volunteer and washed three times in Dulbecco's PBS (PBS, Sigma). A 10% suspension of red blood cells was combined with peptide in a final volume of 200 µl in PBS. PBS was used as a negative control; 0.2% Triton X-100 was used as a positive control. After 30 min incubation, the mixture was  
20 centrifuged 10 min at 3000 rpm in a Beckman TJ6 benchtop centrifuge. Supernatant was carefully transferred to a flat bottom polystyrene 96 well microtiter plate and read at 540 nm. Percent hemolysis was calculated using the formula:  $(A_{\text{Sample}} - A_{\text{blank}}) / (A_{\text{Triton}} - A_{\text{blank}}) \times 100$ .

#### 25 **Example 6: Ovine *In Vivo* Pulmonary Infection Model**

Fifteen lambs were randomized to four groups and were lightly sedated with 10 mg xylazine. A small plastic tube was inserted into the oral cavity and served as a speculum. A bronchoscope was inserted into the tube and lidocaine (0.5 ml of 2% lidocaine hydrochloride) was administered at the larynx. The tip of the bronchoscope  
30 was placed into the dorsum of the caudal portion of the right cranial lobe (pulmonary deposition site) in each animal for deposition of bacterial suspension and treatment

inocula). After inoculation, the effects of xylazine were reversed by intravenous injection of 100 mg tolazoline HCl. Lambs in Group I (2 lambs) and Group II (3 lambs) received 10 ml PBS/PS. Lambs in Group III (5 lambs) and Group IV (5 lambs) received 10 ml PBS/PS and  $2.5 \times 10^8$  CFU/ml *P. haemolytica* in PBS/PS, respectively.

5

At 24 hr post inoculation (PI), lambs in Group I and Group III received an additional 10 ml PBS/PS at the same deposition site. Lambs in Group II received 10 ml PBS/PS with 100 ug SMAP/ml and Group IV received 10 ml PBS/PS with 50 ug SMAP29/ml. At 48 h PI, all lambs were euthanized with pentobarbital and  
10 exsanguinated. At necropsy, the lungs were evaluated grossly and total lung involvement was calculated as the sum of the consolidated portions of each lobe multiplied by the percent each lobe contributes to the total lung volume as previously described (Brogden, 199). The pulmonary inoculation and treatment site was lavaged with 20 ml of PBS. The bronchoalveolar lavage fluids (BALF) were collected for quantitative culture and  
15 determination of total cell count. Tissue was then taken for quantitative bacteriological culture and isolated bacterial species were identified by conventional methodology. All *P. haemolytica* isolates were serotyped to confirm their identity as the original organism.

Pieces of tissue were also taken from the pulmonary deposition site and fixed in  
20 10% neutral buffered formalin solution, dehydrated and cleared, embedded in paraffin, sectioned, and stained with Giemsa and hematoxylin and eosin stains. Lung sections were scored histopathologically in terms of cellular infiltration (lymphocytic and/or neutrophilic), necrosis and collapse as previously described (Brogden, 199). The maximum score was 4 for each category.

25

#### Example 7: Statistics

Statistical differences among gross pulmonary lesions, histopathological lesions, pulmonary cells and *P. haemolytica* in BAL fluids, and *P. haemolytica* in consolidated pulmonary tissue were determined with the Sigmastat 2.03 and Sigmaplot 4.01 (SPSS  
30 Inc., Chicago, IL).

**Example 8: Results**

The following tables summarize the results of experiments described in the preceding section, as do the attached figures and related figure descriptions. These tables exemplify the efficacy of the disclosed peptides and reflect specific peptides which

5 applicants feel have maximum effectiveness.

TABLE 8A. INHIBITION OF *P. AERUGINOSA* PAO1 AND *E. COLI* DH5α BY CAP18 PEPTIDES <sup>1</sup>

Peptide	Sequence	Ionic Strength (mM)	<i>P. aeruginosa</i> PAO1			<i>E. coli</i> DH5α		
			μg/ml ± sem	EC <sub>50</sub> μM ± sem	N	μg/ml ± sem	EC <sub>50</sub> μM ± sem	N
CAP18	GLRKRLRKFRNKIKEKLLKKIG	25	0.98 ± 0.24	0.22 ± 0.05	(6)	0.69 ± 0.08	0.15 ± 0.02	(4)
	QKIQGLLPKLPAPRTDY	175	0.50 ± 0.10	0.11 ± 0.02	(4)			
CAP18-21a	GLRKRLRKFRNKIKEKLLKKIG	25	0.58 ± 0.09	0.22 ± 0.03	(3)	0.48 ± 0.03	0.19 ± 0.01	(3)
		175	0.54 ± 0.10	0.21 ± 0.04	(3)			
CAP18-18	KRLRKFRNKIKEKLLKKIG	25	1.05 ± 0.37	0.46 ± 0.16	(3)	0.47 ± 0.06	0.20 ± 0.03	(3)
		175	1.59 ± 0.29	0.69 ± 0.13	(3)			
CAP18-22	RKRLRKFRNKIKEKLLKKIGQKI	25	1.38 ± 0.49	0.49 ± 0.17	(4)	1.47 ± 0.44	0.52 ± 0.16	(3)
		175	0.88 ± 0.12	0.31 ± 0.04	(4)			
CAP18-19	LRKFRNKIKEKLLKKIGQKI	25	1.65 ± 0.27	0.70 ± 0.11	(3)	2.27 ± 0.30	0.96 ± 0.13	(4)
		175	3.06 ± 0.36	1.29 ± 0.15	(3)			
CAP18-21b	LRKFRNKIKEKLLKKIGQKIQQ	25	1.83 ± 0.51	0.72 ± 0.20	(4)	2.99 ± 0.33	1.17 ± 0.13	(3)
		175	3.45 ± 0.02	1.35 ± 0.01	(2)			
CAP18-15a	RKFRNKIKEKLLKKIG	25	1.54 ± 0.43	0.81 ± 0.23	(3)	0.46 ± 0.12	0.24 ± 0.07	(3)
		175	>20	>10	(3)			
CAP18-15b	KIKEKLLKKIGQKIQQ	25	10.3 ± 0.3	5.92 ± 0.20	(3)	7.23 ± 1.81	4.15 ± 1.04	(4)
		175	>80	>45	(3)			
CAP18-17	KIKEKLLKKIGQKIQGLL	25	2.33 ± 0.35	1.18 ± 0.18	(4)	2.51 ± 0.38	1.27 ± 0.20	(4)
		175	>25	>13	(3)			

<sup>1</sup> EC<sub>50</sub>s were measured in the luminescence assay, performed at 25 mM and 175 mM ionic strength with PAO1 and 25mM with DH5α. Data are average ± Sem.



**TABLE 8B. INHIBITION OF *P. AERUGINOSA* PAO1 BY SMAP29 PEPTIDES <sup>2</sup>**

Peptide	Sequence	Ionic Streng th (mM)	EC <sub>50</sub> <i>P. aeruginosa</i> PAO1		
			μg/ml ± sem	μM ± sem	N
SMAP29	RGLRRLGRKIAHGVKKYGPT VLRIIRIAG	25	0.17 ± 0.02	0.05 ± 0.01	(8)
		175	0.21 ± 0.03	0.06 ± 0.01	(7)
SMAP29-18	RGLRRLGRKIAHGVKKYG	25	0.43 ± 0.08	0.21 ± 0.04	(7)
		175	0.76 ± 0.09	0.37 ± 0.05	(7)
SMAP29-21	KIAHGVKKYGPTVLRIIRIAG	25	1.09 ± 0.16	0.48 ± 0.07	(5)
		175	16.6 ± 1.41	7.25 ± 0.62	(3)
SMAP29-20	LGRKIAHGVKKYGPTVLRII	25	0.83 ± 0.10	0.37 ± 0.05	(5)
		175	27.3 ± 3.97	12.3 ± 1.79	(5)
SMAP29- 18AA	RGLRALGRKIAHGVKAYG	25	0.62 ± 0.13	0.32 ± 0.07	(3)
		175	7.85 ± 1.16	4.08 ± 0.87	(3)
SMAP29X	KNLRRIRKIIHIIKKYGPTILR IIRIIG-NH2	25	2.74 ± 0.65	0.78 ± 0.19	(4)
		175	4.51 ± 0.24	1.29 ± 0.07	(3)
SMAP29X- 18A	KNLRRIRKIIHIIKKYG-NH2	25	0.22 ± 0.00	0.10 ± 0.00	(3)
		175	0.49 ± 0.02	0.22 ± 0.01	(3)
SMAP29X- 18B	KNIRRIIRKIIHIIKKYG-NH2	25	0.16 ± 0.01	0.07 ± 0.00	(3)
		175	0.54 ± 0.05	0.24 ± 0.02	(3)
SMAP29X- 18C	KNIRRIIRKIIHIIKKYG	25	0.24 ± 0.01	0.11 ± 0.00	(3)
		175	0.61 ± 0.12	0.27 ± 0.05	(3)
SMAP29X- 18D	KNLRRIRKIIHIIKKYG	25	0.29 ± 0.01	0.13 ± 0.01	(3)
		175	0.58 ± 0.05	0.26 ± 0.02	(3)
SMAP29X- 16A	NLRRIIRKIIHIIKKY	25	0.90 ± 0.54	0.43 ± 0.26	(3)
		175	1.78 ± 0.26	0.86 ± 0.12	(3)
SMAP29X- 16B	NLRRIIRKIIHIIKKY-NH2	25	0.75	0.36	(1)
		175	3.78	1.82	(1)
SMAP29X- 16C	NIRRIIRKIIHIIKKY-NH2	25	0.65	0.31	(1)
		175	2.21	1.07	(1)
SMAP29X- 16D	NIRRIIRKIIHIIKKY	25	0.70 ± 0.36	0.34 ± 0.17	(3)
		175	3.64 ± 1.75	1.75 ± 0.84	(3)
SMAP29X- 14A	LRRIIRKIIHIIKK-NH2	25	0.23 ± 0.03	0.13 ± 0.02	(3)
		175	1.59 ± 0.22	0.88 ± 0.12	(3)
SMAP29X- 14B	LRRIIRKIIHIIKK	25	0.55 ± 0.03	0.30 ± 0.02	(3)
		175	5.55 ± 0.42	3.08 ± 0.23	(3)
SMAP29X- 14C	IRRIIRKIIHIIKK-NH2	25	0.29 ± 0.04	0.16 ± 0.02	(3)
		175	1.21 ± 0.05	0.67 ± 0.03	(3)

<sup>2</sup> EC<sub>50</sub>S were measured in the luminescence assay, performed at 25 mM and 175 mM ionic strength. Data are average ± sem.

Peptide	Sequence	Ionic Strength (mM)	EC <sub>50</sub> <i>P. aeruginosa</i> PAO1		
			μg/ml ± sem	μM ± sem	N
SMAP29X-14D	IRRIIRKIIHIIKK	25	0.65 ± 0.06	0.36 ± 0.03	(3)
		175	7.11 ± 2.40	3.95 ± 1.34	(3)
SMAP29X-13A	LRRIIRKIIHIIK-NH2	25	0.27 ± 0.03	0.16 ± 0.02	(3)
		175	2.23 ± 0.41	1.34 ± 0.25	(3)
SMAP29X-13B	RRIIRKIIHIIKK-NH2	25	0.22 ± 0.02	0.13 ± 0.01	(3)
		175	1.51 ± 0.09	0.90 ± 0.06	(3)
SMAP29X-12	RRIIRKIIHIIK-NH2	25	0.31 ± 0.01	0.20 ± 0.01	(3)
		175	3.83 ± 0.22	2.46 ± 0.14	(3)
SMAP29X-11A	RRIIRKIIHII-NH2	25	0.38 ± 0.04	0.27 ± 0.03	(3)
		175	4.92 ± 0.43	3.44 ± 0.30	(3)
SMAP29X-11B	RIIRKIIHIIK-NH2	25	0.46 ± 0.02	0.33 ± 0.01	(3)
		175	10.42 ± 0.26	7.43 ± 0.19	(3)
SMAP29X-10	RIIRKIIHII-NH2	25	0.51 ± 0.00	0.40 ± 0.00	(3)
		175	18.67 ± 0.97	14.66 ± 0.77	(3)

**Table 9**  
**Relative Time for Killing of *P. aeruginosa* PAO1 by CAP18 and SMAP29 Peptides<sup>1</sup>**

Peptide	Peptide ( $\mu\text{g/ml}$ )	Half-time of Killing (min. avg. $\pm$ sem)
CAP18	10	4.0 + 1.4
CAP18-21A	6	9.2 $\pm$ 3.4
CAP18-18	10	19.5 $\pm$ 8.2
CAP18-22	14	9.8 $\pm$ 2.5
CAP18-19	16	8.0 $\pm$ 1.6
SMAP29	2	2.3 $\pm$ 0.8
SMAP29X-18D	5 <sup>a</sup>	2.9
SMAP29X-14A	4 <sup>b</sup>	20.9

5                   <sup>1</sup> Peptides were used at a concentration of ten times their EC<sub>50</sub>, except where  
noted, with *P. aeruginosa* at 25 mM ionic strength (Tables 8A and 8B). Bacteria were  
incubated with peptide, and luminescence was measured at intervals. The half-time of  
killing is that time at which luminescence was decreased to half its original value. <sup>a</sup>The  
10 concentration of SMAP29X-18D was 15.7 X the EC<sub>50</sub>. <sup>b</sup>The concentration of  
SMAP29X-14A was 17.4 X the EC<sub>50</sub>.

**TABLE 10A**  
**MULTIPLY DRUG-RESISTANT STRAINS OF *P. AERUGINOSA* FROM CF**  
**PATIENTS ARE INHIBITED BY CAP18 PEPTIDES<sup>3</sup>**  
**( MIC,  $\mu$ g/ml)**

Strain #	ID	CAP18	CAP18-22	CAP18-18
85BM	P.a	2	4	1 6
45BN	P.a	1	1	2
83BN	P.a	1	4	1 6
50BK	P.a	4	2	8
18BA	P.a	4	2	2
75BN	P.a	8	3 2	>3 2
13BK	M P.a	4	2	1 6
7BN	M P.a	2	4	1 6
1BH	M P.a	1	4	8
26BL	M P.a	8	1 6	>3 2
38BJ	M P.a	1	0.5	1
68BM	M P.a	4	8	3 2

5

P.a: Nonmucoid

M P.a: Mucoid

These data were provided by Dr. Lisa Saiman, Dept. of Pediatrics,  
Columbia University, New York, NY.

<sup>3</sup> Determined by Microbroth Dilution Assay.

TABLE 10B  
MULTIPLY DRUG-RESISTANT STRAINS OF *P. AERUGINOSA* FROM CF PATIENTS ARE INHIBITED BY SMAP29 PEPTIDES<sup>4</sup>  
(MIC, µg/ml)

Strain#	ID	SMAP29	SMAP29-18	SMAP29X-14A	SMAP29X-14B/D	SMAP29X-18A	SMAP29X-18C	SMAP29X-18D
18BA	P.a	1	8	2	>32	8	4	8
73BI	P.a	2	>32	2	>32	8	16	8
3BJ	P.a	0.5	4	2	>32	4	8	4
23BJ	P.a	8	8	16	>32	>32	32	16
38BK	P.a	1	32	2	>32	8	8	8
50BK	P.a	2	8	2	>32	4	4	4
1BL	P.a	2	>32	16	>32	16	16	16
26BL	M P.a	2	>32	4	>32	16	16	8
41BL	M P.a	0.5	16	2	>32	2	4	4
84BL	M P.a	1	>32	4	>32	4	4	8
68BM	M P.a	2	32	1	>32	8	8	8
82BM	M P.a	1	32	0.06	>32	4	4	4
1BH	M P.a	0.5	4	2	>32	2	2	2
13BH	M P.a	0.13	1	0.5	>32	0.5	0.06	0.12
32BH	M P.a	1	32	2	>32	4	4	4
38BJ	M P.a	0.5	32	4	>32	>32	2	16
95BJ	M P.a	0.5	16	1	>32	2	1	2
13BK	M P.a	1	16	2	>32	4	8	4
57BK	M P.a	0.5	32	4	>32	4	4	4

P.a: Nonmucoid  
M P.a: Mucoid

These data were provided by Dr. Lisa Saiman, Dept. of Pediatrics, Columbia University, New York, NY.

<sup>4</sup> Determined by Microbroth Dilution Assay.

TABLE 11  
INHIBITION OF OTHER MULTIPLY DRUG-RESISTANT BACTERIA FROM  
CF PATIENTS BY SMAP29 AND CAP18 PEPTIDES<sup>5</sup>  
(MIC, µg/ml)

Strain #	API-ID	SMAP29	CAP18
29AX	A. x	1	2
34AX	A. x	3 2	>3 2
39BB	A. x	3 2	>3 2
61AW	A. x	3 2	3 2
62AZ	A. x	3 2	1 6
33BI	B. c	>32	>3 2
45BI	B. c	>32	>3 2
54BK	B. c	>32	>3 2
90BD	B. c	>32	>3 2
96BD	B. c	>32	>3 2
59BH	X. m	2	4
77BJ	X. m	4	1 6
81BG	X. m	4	1 6
85BG	X. m	8	1 6
100BC	X. m	1	2

A. x: *Alcaligenes xylosoxidans*  
B. c: *Burkholderia cepacia*  
X. m: *Stenotrophomonas maltophilia*  
These data were provided by Dr. Lisa Saiman, Dept. of Pediatrics,  
Columbia University, New York, NY.

<sup>5</sup> Determined by Microbroth Dilution Assay.



**TABLE 12**  
**VANCOMYCIN AND METHICILLIN-RESISTANT GRAM-POSITIVES ARE**  
**INHIBITED BY SMAP29 PEPTIDES<sup>6</sup>**  
**(MIC, µG/ML)**

Organism	Phenotype/ Genotype	Smap29	Smap29X-18D	Smap29X-14A
E.f 142-1498-1	Van A	1 6	3 2	>6 4
E.f 142-6873-3	Van B	3 2	6 4	>6 4
E.fm 110-2733-2	Van B	4	1 6	1 6
E.fm 110-6802-2	Van A	4	8	8
E.fm 115-7485-1	Susc.	4	1 6	1 6
S.e Dade 352	MecA+	8	8	8
S.e Dade 354	MecA+	4	4	4
S.e Dade 370	MecA-	4	4	8
S.e Dade 371	MecA-	4	4	8
S.a ATCC 25932	MecA-	8	1 6	6 4
S.a ATCC 29213	MecA-	8	1 6	3 2
S.a Dade 377	MecA+	8	1 6	6 4
S.a Dade 382	MecA+	8	1 6	3 2
S.a Mu 3	Visa	1 6	3 2	6 4
S.a Mu 50	Visa	1 6	3 2	3 2

E.f: Enterococcus faecalis

Mec A+: Methicillin-resistant

E.fm: Enterococcus faecium

Mec A-: Methicillin-sensitive

S.a: Staphylococcus aureus

Visa: Vancomycin intermediately-resistant

*S.aureus*

S.e: Staphylococcus epidermidis Van A,B: Vancomycin-resistant

These data were provided by Dr. Patrica Winokur, Dept. of Internal Medicine, U of I,  
 Iowa City, IA.

<sup>6</sup> Determined by Microbroth Dilution Assay.

**TABLE 13**  
**ANTIPSEUDOMONAL AND ANTISTAPHYLOCOCCAL ACTIVITIES OF**  
**CAP18 AND SMAP29 PEPTIDES ASSESSED BY A RADIAL DIFFUSION**  
**ASSAY (MIC, µg/ml)**

5

	<i>E. coli</i>		<i>P. aeruginosa</i>		MRSA		<i>S. aureus</i>	
	[NaCl]		[NaCl]		[NaCl]		[NaCl]	
	Low	High	Low	High	Low	High	Low	High
SMAP29 and Related Peptides								
SMAP29	0.6	0.4	1.1	0.6	7.1	3.3	1.0	1.4
SMAP29-18	1.9	0.4	1.0	9.9	8.7	>250	2.4	24.5
SMAP29X	1.5	0.8	2.6	1.0	6.2	1.0	1.6	1.5
SMAP29X-18D	0.8	0.5	1.1	0.5	8.7	0.8	2.0	1.4
SMAP29X-14A	2.9	0.9	2.0	2.7	7.6	>79.1	1.1	1.9
CAP18 and Related Peptides								
CAP18	0.2	0.4	0.2	1.6	8.2	6.1	1.8	2.8
CAP18-22	0.1	1.7	0.7	7.6	8.1	>79.1	7.9	>79.1
CAP18-21a	2.3	0.4	0.7	0.9	19.4	>250	1.8	2.2
CAP18-18	2.8	1.5	2.1	3.2	14.9	>250	9.7	29.0
CAP18-15a	0.2	0.7	1.5	5.3	>250	>250	19.5	>250
CAP18-21b	2.3	0.5	2.9	23.5	10.3	>250	16.3	>250
CAP18-19	0.4	18.2	1.0	26.7	4.5	>79.1	6.0	>79.1
CAP18-17	9.7	1.9	5.1	>250	>250	>250	>250	>250
CAP18-15b	1.6	>250	8.7	>250	>250	>250	>250	>250

10 Assays were performed in low salt (10mM sodium phosphate buffer, pH 7.4) and high salt underlays (10mM sodium phosphate buffer, +100 mM NaCl, pH 7.4). The bacterial strains were *E. coli* ML-35p, *P. aeruginosa* MR 3007, *S. aureus* 930918-3 and MRSA, ATCC 33591. These data were provided by Dr. Robert I. Lehrer, Dept. of Medicine, UCLA, Los Angeles, CA.

TABLE 14  
CONCENTRATION OF *PASTEURELLA HAEMOLYTICA*<sup>1</sup> IN  
BRONCHOAVEOLAR LAVAGE FLUID (BALF) AND CONSOLIDATED LUNG  
TISSUE (CLT) OF INFECTED SHEEP

Group	Log <sub>10</sub> CFU/ml of BALF	Log <sub>10</sub> CFU/gm of clt
1 Control: PBS/PS <sup>4</sup>	0	PBS/PND
2 Control: PBS/PS/SMAP29	0	ND
3 <i>P. haemolytica</i>	2.9 ± 0.7 <sup>2</sup>	5.1 ± 1.0 <sup>2</sup>
4 <i>P. haemolytica</i> + SMAP29	1.6 ± 0.4 <sup>3</sup>	3.7 ± 0.9 <sup>3</sup>

- 10
- 1

2

3

4

These data were provided by Dr. Kim Brogden, NADC, ARS, USDA, Ames, IA.
- Serotype A1, strain 82-25

Mean CFU in 4 of 5 lambs

Mean CFU in 3 of 5 lambs

Phosphate Buffered Saline pH 7.4/Ovine Pulmonary Surfactant

Example 9

Susceptibility of a variety of bacterial strains to the disclosed peptides under either normal, high and low salinity were evaluated. Bacterial strains *E. coli* ML-35p, *P. aeruginosa* MR 3007, *S. aureus* 930918-3 and MRSA ATCC 33591 MICs were determined over a broader range of salt concentration, i.e. 0 to 175 mM. The susceptibility of several additional strains of bacteria were also evaluated by the Radial Immunodiffusion Assay including *E. coli* ATCC 33780 (Serotype 0111), *E. coli* DH-Salpha, *P. aeruginosa* PAO1, *N. gonorrhoeae*, *B. cepacia* ATCC 25416 and *L. monocytogenes* EGD. Results of these assays are set forth in Tables 15-24.

**Example 10**

The Smap28 peptide (SEQ ID NO:27), has been is the the physiologically relevant form of the sheep myeloid antimicrobial protein Smap29. The peptide was  
5 purified from sheep leukocytes, and its structure determined by mass spectrometry analysis and Edman degradation. The structure of the Smap28 peptide is represented by two interconvertable conformers of the "helix-hinge-helix" motif. It is believed that the the N- and C-termini of Smap 29 constitute distinct LPS-binding domains, the primary means by which these peptides recognize Gram-negative bacteria (FIGS. 7 & 8).

10

Activities of the peptide were evaluated towards *E. coli* ML-35p, *P. aeruginosa* MR 3007, *L. monocytogenes* EGD and MRSA ATCC 33591 (See Tables 15-17). Activity of the native peptide was determined to be indistinguishable from the synthetic version of Smap28 (FIGS. 9 and 10). Activity was concentration dependent but efficacy  
15 of the native and synthetic proteins was determined to be statistically the same.

Table 15: <i>E. coli</i> ML-35p, MIC( $\mu$ g/ml) in underlays of low, normal and high salinity						
Seq. ID	Primary Structure	0 m M NaCl	100 mM NaCl	175 mM NaCl		
2	RGLRR LGRKI AHGVK KYGPT VLRII RIAG	0.54 $\pm$ 0.15	0.35 $\pm$ 0.05	0.32 $\pm$ 0.08		
3	KI AHGVK KYGPT VLRII RIAG	0.20 $\pm$ 0.09	6.21 $\pm$ 0.62	>79.1		
4	LGRKI AHGVK KYGPT VLRII	0.20 $\pm$ 0.08	21.97 $\pm$ 0.63	>79.1		
5	RGLRR LGRKI AHGVK KYG	0.44 $\pm$ 0.15	5.87 $\pm$ 0.52	58.6 $\pm$ 11.79		
8	KNLRR IIRKI IHIK KYG	0.45 $\pm$ 0.10	0.38 $\pm$ 0.02	0.41 $\pm$ 0.06		
11	LRR IIRKI IHIK K-NH <sub>2</sub>	0.63 $\pm$ 0.05	5.87 $\pm$ 0.04	49.2 $\pm$ 15.2		
12	LRR IIRKI IHIK K	0.68 $\pm$ 0.07	15.9 $\pm$ 5.0	>79.1		
13	IRR IIRKI IHIK K-NH <sub>2</sub>	n.t.	n.t.	n.t.		
15	LRR IIRKI IHIK-NH <sub>2</sub>	0.64 $\pm$ 0.15	2.24 $\pm$ 0.32	12.6 $\pm$ 2.8		
17	RR IIRKI IHIK-NH <sub>2</sub>	0.29 $\pm$ 0.06	17.6 $\pm$ 5.73	>79.1		
18	GLRKR LRKER NKIKE KLKKI GQKIQ GLLPK LAPRT DY	2.8	1.5	n.t.		
19	GLRKR LRKER NKIKE KLKKI G	2.3	0.4	n.t.		
20	KR LRKER NKIKE KLKKI G	2.3	0.5	n.t.		
21	RKR LRKER NKIKE KLKKI GQKI	0.2	0.4	n.t.		
22	LRKER NKIKE KLKKI GQKI	0.1	1.7	n.t.		
23	LRKER NKIKE KLKKI GQKIQ G	0.2	0.7	n.t.		
24	RKFR NKIKE KLKKI G	9.7	1.9	n.t.		
25	KIKE KLKKI GQKIQ G	1.6	>250	n.t.		
26	KIKE KLKKI GQKIQ GLL	0.4	18.2	n.t.		
27	RGLRR LGRKI AHGVK KYGPT VLRII RIA-NH <sub>2</sub>	0.37 $\pm$ 0.09	0.34 $\pm$ 0.02	0.42 $\pm$ 0.17		
28	KNLRR IIRKI IHIK KYGPT ILRII RIIG-NH <sub>2</sub>	0.16 $\pm$ 0.03	0.52 $\pm$ 0.12	0.58 $\pm$ 0.12		

Table 16: <i>E. coli</i> ATCC 33780 (Serotype 0111) MIC (µg/mL) in underlays of low, normal and high salinity						
Seq. #	Primary Structure		0 m M NaCl	100 mM NaCl	175 mM NaCl	
2	RGLRR	LGRKI AHGVK KYGPT VLRII RIAG	0.46 ± 0.09	0.28 ± 0.01	0.26 ±	
3		KI AHGVK KYGPT VLRII RIAG	0.53 ± 0.24	6.80 ± 0.60	73.2 ± 5.84	
4		LGRKI AHGVK KYGPT VLRII	0.82 ± 0.22	20.8 ± 1.46	>79.1	
5	RGLRR	LGRKI AHGVK KYG	2.98 ± 1.39	9.74 ± 6.03	38.0 ± 20.5	
27	RGLRR	LGRKI AHGVK KYGPT VLRII RIA-NH <sub>2</sub>	0.61 ± 0.15	0.36 ± 0.04	0.36 ± 0.04	



Table 17: <i>E. coli</i> DH-5 $\alpha$ , MIC( $\mu$ g/mL) in underlays of low, normal and high salinity						
Seq. ID	Primary Structure		0 m M NaCl	100 mM NaCl	175 mM NaCl	
2	RGLRR	LGRKI AHGVK KYGPT VLRII RIAG	0.55 $\pm$ 0.15	0.32 $\pm$ 0.04	0.31 $\pm$ 0.08	
3		KI AHGVK KYGPT VLRII RIAG	0.42 $\pm$ 0.10	22.91 $\pm$ 12.01	>79.1	
4		LGRKI AHGVK KYGPT VLRII	0.76 $\pm$ 0.12	47.79 $\pm$ 20.40	>79.1	
5	RGLRR	LGRKI AHGVK KYG	1.15 $\pm$ 0.16	15.63 $\pm$ 4.46	60.03 $\pm$ 13.2	
8	KNLRR	IIRKI IHIK KYG	0.26 $\pm$ 0.05	0.31 $\pm$ 0.06	0.35 $\pm$ 0.09	
11	LRR	IIRKI IHIK K-NH <sub>2</sub>	0.85 $\pm$ 0.04	1.91 $\pm$ 0.40	5.92 $\pm$ 1.02	
12	LRR	IIRKI IHIK K	1.36 $\pm$ 0.19	4.93 $\pm$ 1.41	19.1 $\pm$ 5.0	
13	IRR	IIRKI IHIK K-NH <sub>2</sub>	n.t.	n.t.	n.t.	
15	LRR	IIRKI IHIK-NH <sub>2</sub>	0.82 $\pm$ 0.12	1.84 $\pm$ 0.36	5.35 $\pm$ 1.87	
17	RR	IIRKI IHIK-NH <sub>2</sub>	0.51 $\pm$ 0.09	6.19 $\pm$ 1.67	46.1 $\pm$ 16.1	
18	GLRKR	LRKFR NKIKE KLKKI GQKIQ GLLPK LAPRT DY	0.84	0.91	n.t.	
27	RGLRR	LGRKI AHGVK KYGPT VLRII RIA-NH <sub>2</sub>	0.47 $\pm$ 0.11	0.30 $\pm$ 0.04	0.37 $\pm$ 0.17	
28	KNLRR	IIRKI IHIK KYGPT ILRII RIIG-NH <sub>2</sub>	0.0042 $\pm$ 0.0027	0.27 $\pm$ 0.05	0.40 $\pm$ 0.09	

Table 18: *P. aeruginosa* MR-3007 MIC( $\mu\text{g/ml}$ ) in underlays of low, normal and high salinity

Seq. #	Structure	0 m M NaCl	100 mM NaCl	175 mM NaCl
2	RGLRR LGRKI AHGVK KYGPT VLRII RIAG	0.60 $\pm$ 0.07	0.60 $\pm$ 0.05	0.50 $\pm$ 0.05
3	KI AHGVK KYGPT VLRII RIAG	0.61 $\pm$ 0.12	15.8 $\pm$ 0.62	30.1 $\pm$ 0.9
4	LGRKI AHGVK KYGPT VLRII	0.64 $\pm$ 0.16	44.5 $\pm$ 0.63	>79.1
5	RGLRR LGRKI AHGVK KYG	1.14 $\pm$ 0.29	8.22 $\pm$ 0.52	13.2 $\pm$ 4.2
8	KNLRR IIRKI IHIK KYG	0.67 $\pm$ 0.12	0.57 $\pm$ 0.02	0.59 $\pm$ 0.05
11	LRR IIRKI IHIK K-NH <sub>2</sub>	0.92 $\pm$ 0.06	4.36 $\pm$ 0.04	20.1 $\pm$ 6.9
12	LRR IIRKI IHIK K	0.98 $\pm$ 0.08	9.38 $\pm$ 5.0	46.3 $\pm$ 9.1
13	IRR IIRKI IHIK K-NH <sub>2</sub>	0.84 $\pm$ 0.06	3.99 $\pm$	19.7 $\pm$ 2.2
15	LRR IIRKI IHIK-NH <sub>2</sub>	0.96 $\pm$ 0.08	3.07 $\pm$ 0.32	9.14 $\pm$ 4.81
17	RR IIRKI IHIK-NH <sub>2</sub>	0.72 $\pm$ 0.15	44.6 $\pm$ 5.73	>79.1
18	GLRKR LRKFR NKIKE KLKKI GQKIQ GLLPK LAPRT DY	2.1	3.2	n.t.
19	GLRKR LRKFR NKIKE KLKKI G	0.7	0.9	n.t.
20	KR LRKFR NKIKE KLKKI G	2.9	23.5	n.t.
21	RKR LRKFR NKIKE KLKKI GQKI	0.2	1.6	n.t.
22	LRKFR NKIKE KLKKI GQKI	0.7	7.6	n.t.
23	LRKFR NKIKE KLKKI GQKIQ G	1.5	5.3	n.t.
24	RKFR NKIKE KLKKI G	5.1	>250	n.t.
25	KIKE KLKKI GQKIQ G	8.7	>250	n.t.
26	KIKE KLKKI GQKIQ GLL	1.0	26.7	n.t.
28	KNLRR IIRKI IHIK KYGPT ILRII RIIG-NH <sub>2</sub>	0.34 $\pm$ 0.22	0.51 $\pm$ 0.04	0.49 $\pm$ 0.06

Table 19: <i>P. aeruginosa</i> PA-01 MIC( $\mu\text{g/mL}$ ) in underlays of low, normal and high salinity					
Seq. #	Structure	0 m M NaCl	100 mM NaCl	175 mM NaCl	
2	RGLRR LGRKI AHGVK KYGPT VLRII RIAG	0.78 $\pm$ 0.01	0.59 $\pm$ 0.03	0.51 $\pm$ 0.03	
3	KI AHGVK KYGPT VLRII RIAG	0.44 $\pm$ 0.06	13.6 $\pm$ 7.56	32.2 $\pm$ 0.7	
4	LGRKI AHGVK KYGPT VLRII	0.47 $\pm$ 0.10	39.0 $\pm$ 20.1	>79.1	
5	RGLRR LGRKI AHGVK KYG	0.28 $\pm$ 0.04	1.72 $\pm$ 0.6	5.40 $\pm$ 1.25	
8	KNLRR IIRKI IHIK KYG	0.83 $\pm$ 0.05	0.70 $\pm$ 0.02	0.82 $\pm$ 0.05	
11	LRR IIRKI IHIK K-NH <sub>2</sub>	0.96 $\pm$ 0.05	2.28 $\pm$ 1.02	6.51 $\pm$ 1.02	
12	LRR IIRKI IHIK K	0.98 $\pm$ 0.05	4.19 $\pm$ 1.63	11.9 $\pm$ 4.7	
13	IRR IIRKI IHIK K-NH <sub>2</sub>	1.03 $\pm$ 0.08	1.98 $\pm$ 0.74	7.45 $\pm$ 0.87	
15	LRR IIRKI IHIK-NH <sub>2</sub>	1.07 $\pm$ 0.19	4.86 $\pm$ 2.69	4.66 $\pm$ 1.57	
17	RR IIRKI IHIK-NH <sub>2</sub>	0.69 $\pm$ 0.08	7.09 $\pm$ 2.80	17.2 $\pm$ 4.6	
18	GLRKR LRKFR NKIKE KLKKI GQKIQ GLLPK LAPRT DY	0.88	2.77	n.t.	
28	KNLRR IIRKI IHIK KYGPT ILRII RIIG-NH <sub>2</sub>	0.53 $\pm$ 0.14	0.70 $\pm$ 0.09	1.17 $\pm$ 0.23	

Table 21: <i>N. gonorrhoeae</i> -done only at 100mM NaCl				
Seq. #	Structure	FA 19 (serum-resistant)	F 62 (serum-sensitive)	
2	RGLRR LGRKI AHGVK KYGPT VLRII RIAG	0.2	0.3	
5	RGLRR LGRKI AHGVK KYG	56.9	20.6	
8	KNLRR IIRKI IHIK KYG	0.2	0.5	
11	LRR IIRKI IHIK K-NH <sub>2</sub>	0.9	0.8	
18	GLRKR LRKFR NKIKE KLKKI GQKIQ GLLPK LAPRT DY	0.6	0.7	
19	GLRKR LRKFR NKIKE KLKKI G	>79.1	0.6	
20	KR LRKFR NKIKE KLKKI G	>250	>250	
21	RKR LRKFR NKIKE KLKKI GQKI	>250	22.0	
22	LRKFR NKIKE KLKKI GQKI	>250	>79.1	
23	LRKFR NKIKE KLKKI GQKIQ G	>250	>79.1	
24	RKFR NKIKE KLKKI G	>250	>250	
25	KIKE KLKKI GQKIQ G	>250	>79.1	
26	KIKE KLKKI GQKIQ GLL	>250	>79.1	
28	KNLRR IIRKI IHIK KYGPT ILRII RIIG-NH <sub>2</sub>	0.1	0.1	

Table 22: <i>S. aureus</i> 930918-3 MIC( $\mu\text{g/mL}$ ) in underlays of low, normal and high salinity						
Seq. #	Structure	0 m M NaCl	100 mM NaCl	175 mM NaCl		
2	RGLRR LGRKI AHGVK KYGPT VLRII RIAG	1.01 $\pm$ 0.20	0.82 $\pm$ 0.08	1.62 $\pm$ 0.35		
3	KI AHGVK KYGPT VLRII RIAG	2.57 $\pm$ 0.23	26.7 $\pm$ 2.5	>79.1		
4	LGRKI AHGVK KYGPT VLRII	6.97 $\pm$ 0.59	>79.1	>79.1		
5	RGLRR LGRKI AHGVK KYG	6.75 $\pm$ 0.51	>79.1	>79.1		
8	KNLRR IIRKI IHIK KYG	0.84 $\pm$ 0.14	0.79 $\pm$ 0.02	0.86 $\pm$ 0.06		
11	LRR IIRKI IHIK K-NH <sub>2</sub>	1.38 $\pm$ 0.21	4.70 $\pm$ 1.36	16.4 $\pm$ 5.1		
12	LRR IIRKI IHIK K	2.40 $\pm$ 0.17	65.4 $\pm$ 13.7	79.1 $\pm$ 0		
13	IRR IIRKI IHIK K-NH <sub>2</sub>	1.03 $\pm$ 0.03	2.01 $\pm$ 0.06	5.93 $\pm$ 1.12		
15	LRR IIRKI IHIK-NH <sub>2</sub>	1.32 $\pm$ 0.22	3.88 $\pm$ 0.93	6.39 $\pm$ 1.12		
17	RR IIRKI IHIK-NH <sub>2</sub>	2.29 $\pm$ 0.07	>79.1	>79.1		
18	GLRKR LRKFR NKIKE KLKKI GQKIQ GLLPK LAPRT DY	1.8	2.8	n.t.		
19	GLRKR LRKFR NKIKE KLKKI G	1.8	2.2	n.t.		
20	KR LRKFR NKIKE KLKKI G	9.7	29.0	n.t.		
21	RKR LRKFR NKIKE KLKKI GQKI	7.9	>79.1	n.t.		
22	LRKFR NKIKE KLKKI GQKI	6.0	>79.1	n.t.		
23	LRKFR NKIKE KLKKI GQKIQ G	16.3	>250	n.t.		
24	RKFR NKIKE KLKKI G	19.5	>250	n.t.		
25	KIKE KLKKI GQKIQ G	>250	>250	n.t.		
26	KIKE KLKKI GQKIQ GLL	>250	>250	n.t.		
28	KNLRR IIRKI IHIK KYGPT ILRII RIIG-NH <sub>2</sub>	0.55 $\pm$ 0.17	0.82 $\pm$ 0.04	1.14 $\pm$ 0.23		

Table 23: MRSA ATCC 33591 MIC( $\mu\text{g/mL}$ ) in underlays of low, normal and high salinity						
Seq. #	Primary Structure		0 m M NaCl	100 mM NaCl	175 mM NaCl	
2	RGLRR	LGRKI AHGVK KYGPT VLRII RIAG	0.77 $\pm$ 0.08	0.90 $\pm$ 0.17	1.34 $\pm$ 0.23	
3		KI AHGVK KYGPT VLRII RIAG	2.53 $\pm$ 0.07	>79.1	>79.1	
4		LGRKI AHGVK KYGPT VLRII	7.03 $\pm$ 0.66	>79.1	>79.1	
5	RGLRR	LGRKI AHGVK KYG	5.28 $\pm$ 0.49	>79.1	>79.1	
8	KNLRR	IIRKI IHIK KYG	0.77 $\pm$ 0.04	0.75 $\pm$ 0.06	0.89 $\pm$ 0.05	
11	LRR	IIRKI IHIK K-NH <sub>2</sub>	4.42 $\pm$ 1.73	23.6 $\pm$ 1.6	>79.1	
12	LRR	IIRKI IHIK K	11.0 $\pm$ 3.96	>79.1	>79.1	
13	IRR	IIRKI IHIK K-NH <sub>2</sub>	2.39 $\pm$ 0.07	21.6 $\pm$ 0.3	>79.1	
15	LRR	IIRKI IHIK-NH <sub>2</sub>	2.49 $\pm$ 0.27	24.7 $\pm$ 2.5	65.8 $\pm$ 13.3	
17	RR	IIRKI IHIK-NH <sub>2</sub>	11.9 $\pm$ 4.7	>79.1	>79.1	
18	GLRKR	LRKER NKIKE KLKKI GQKIQ GLLPK LAPRT DY	8.2	2.8	n.t.	
19	GLRKR	LRKER NKIKE KLKKI G	19.4	2.2	n.t.	
20	KR	LRKER NKIKE KLKKI G	14.9	29.0	n.t.	
21	RKR	LRKER NKIKE KLKKI GQKI	8.1	>79.1	n.t.	
22		LRKER NKIKE KLKKI GQKI	4.5	>79.1	n.t.	
23		LRKER NKIKE KLKKI GQKIQ G	10.3	>250	n.t.	
24		RKER NKIKE KLKKI G	>250	>250	n.t.	
25		KIKE KLKKI GQKIQ G	>250	>250	n.t.	
26		KIKE KLKKI GQKIQ GLL	>250	>250	n.t.	
28	KNLRR	IIRKI IHIK KYGPT ILRII RIIG-NH <sub>2</sub>	0.32 $\pm$ 0.04	0.81 $\pm$ 0.07	1.57 $\pm$ 0.30	



Table 24: <i>L. monocytogenes</i> in underlays of low, normal and high salinity					
Seq. #	Structure	0 m M NaCl	100 mM NaCl	175 mM NaCl	
2	RGLRR LGRKI AHGVK KYGPT VLRII RIAG	0.5	0.4	n.t.	
5	RGLRR LGRKI AHGVK KYG	1.0	9.7	n.t.	
8	KNLRR IIRKI IHIK KYG	2.1	0.7	n.t.	
11	LRR IIRKI IHIK K-NH <sub>2</sub>	0.7	0.5	n.t.	
18	GLRKR LRKFR NKIKE KLKKI GQKIQ GLLPK LAPRT DY	0.6	0.7	n.t.	
19	GLRKR LRKFR NKIKE KLKKI G	0.3	0.5	n.t.	
20	KR LRKFR NKIKE KLKKI G	2.3	>250	n.t.	
21	RKR LRKFR NKIKE KLKKI GQKI	7.0	30.3	n.t.	
22	LRKFR NKIKE KLKKI GQKI	5.9	>79.1	n.t.	
23	LRKFR NKIKE KLKKI GQKIQ G	5.5	>79.1	n.t.	
24	RKFR NKIKE KLKKI G	24.1	>250	n.t.	
25	KIKE KLKKI GQKIQ G	>250	>250	n.t.	
26	KIKE KLKKI GQKIQ GLL	5.7	>79.1	n.t.	
28	KNLRR IIRKI IHIK KYGPT ILRII RIIG-NH <sub>2</sub>	0.4	0.5	n.t.	

\*\*\*\*\*

5 All of the COMPOSITIONS and/or METHODS disclosed and claimed herein can  
be made and executed without undue experimentation in light of the present disclosure.  
While the compositions and methods of this invention have been described in terms of  
preferred embodiments, it will be apparent to those of skill in the art that variations may  
be applied to the COMPOSITIONS and/or METHODS and in the steps or in the  
sequence of steps of the method described herein without departing from the concept,  
10 spirit and scope of the invention. More specifically, it will be apparent that certain agents  
which are both chemically and physiologically related may be substituted for the agents  
described herein while the same or similar results would be achieved. All such similar  
substitutes and modifications apparent to those skilled in the art are deemed to be within  
the spirit, scope and concept of the invention as defined by the appended claims.

## REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein  
5 by reference.

Agerberth *et al.*, *Proc. Nat'l Acad. Sci. USA*, 92:195-199, 1995.

Bagella *et al.*, *FEBS Lett.*, 376:225-228, 1995.

Baichwal and Sugden, In: *Gene Transfer*, Kucherlapati R, ed., New York, Plenum Press,  
10 117-148, 1986.

Bals *et al.*, *J. Clin. Invest.*, 102:874-880, 1998.

Bals *et al.*, "The peptide antibiotic LL-37/hCAP-18 is expressed in epithelia of the human lung where it has broad antimicrobial activity at the airway surface," *Proc. Nat'l Acad. Sci. USA*, 95:9541-9546, 1998.

15 Bangham *et al.*, *J. Mol. Biol.*, 13: 238-252, 1965.

Bennik *et al.*, "A novel bacteriocin with a YGNGV motif from vegetable-associated *Enterococcus mundtii*: full characterization and interaction with target organisms," *Biochim Biophys Acta*, 1373:47-58, 1998.

Benvenisty and Neshif, *Proc. Nat'l Acad. Sci. USA*, 83:9551-9555, 1986.

20 Berklman *et al.*, *J. Infect Dis.*, 170(2):272-7, 1994.

Boman, "Peptide antibiotics and their role in innate immunity," *Annu. Rev. Immunol.*, 13:61-92, 1995.

Boman, *Annu. Rev. Immunol.*, 13, 61-92, 1995.

Brasseur *et al.*, *Trends in Biochem Soc.*, 22:167-71, 1997.

25 Capaldi *et al.*, *Biochem Biophys Res Commun*, 74(2):425-33, 1977.

Chang *et al.*, *Hepatology*, 14:124A, 1991.

Chen and Okayama, *Mol. Cell Biol.*, 7:2745-2752, 1987.

Chen *et al.*, *FEBS Lett.*, 370:46-52, 1995.

Cleeland and Squires, In: *Lorian, Antibiotics in Laboratory Medicine*, Satterfield (Ed),  
30 Williams & Wilkins, Philadelphia, 1991.

- Coffin, *In: Fields BN, Knipe DM, ed. Virology*. New York: Raven Press, pp. 1437-1500, 1990.
- Couch *et al.*, *Am. Rev. Resp. Dis.*, 88:394-403, 1963.
- Coupar *et al.*, *Gene*, 68:1-10, 1988.
- 5 Davies, *FEMS Microbiology Reviews*, 39:363-371, 1986.
- Deamer and Uster, "Liposome Preparation: Methods and Mechanisms," *In: Liposomes*, M. Ostro (Ed.), 1983
- Dubensky *et al.*, *Proc. Nat'l Acad. Sci. USA*, 81:7529-7533, 1984.
- Eisenberg, D. *et al. J Cell Biochem* 31: 11-7, 1986.8
- 10 Epand *et al.*, *Biopolymers*, 37:319-338, 1995.
- EPO 0273085
- Fechheimer *et al.*, *Proc. Nat'l. Acad. Sci. USA*, 84:8463-8467, 1987.
- Ferkol *et al.*, *FASEB J.*, 7:1081-1091, 1993.
- Frackman *et al.*, *J. Bacteriol.*, 172:5767-5773, 1990.
- 15 Fraley *et al.*, *Proc. Nat'l Acad. Sci. USA*, 76:3348-3352, 1979.
- Freshner, *In: Animal Cell Culture: a Practical Approach* Second Edition, Oxford/New York, IRL Press, Oxford University Press, 1992.
- Friedmann, *Science*, 244:1275-1281, 1989.
- Frohm Nilsson *et al.*, *Infect Immun*, 67(5):2561-6, 1999.
- 20 Gallo *et al.*, *J. Biol. Chem.*, 272:13088-13093, 1997.
- Gallo *et al.*, *J Invest Dermatol.*, 111(5):739-43, 1998.
- Ganz and Lehrer, *Pharmac. Ther.*, 66:191-205, 1995.
- Ganz and Lehrer, *Current Opinions Immun.*, 10:41, 1998.
- Ghosh and Bachhawat, *In: Wu G. and C. Wu (eds.) Liver diseases, targeted diagnosis*
- 25 and therapy using specific receptors and ligands, New York: Marcel Dekker, pp. 87-104, 1991.
- Ghosh-Choudhury *et al.*, *EMBO J.*, 6:1733-1739, 1987.
- Goldman *et al.*, *Cell*, 88:553-560, 1997.
- Gomez-Foix *et al.*, *J. Biol. Chem.*, 267:25129-25134, 1992.
- 30 Gopal, *Mol. Cell Biol.*, 5:1188-1190, 1985.
- Graham and Prevec, *Biotechnology*, 20:363-390, 1992.

- Graham and Prevec, In: E.J. Murray (ed.), *Methods in Molecular Biology: Gene Transfer and Expression Protocol*, Clifton, NJ: Humana Press, 7:109-128, 1991.
- Graham and Van Der Eb, *Virology*, 52:456-467, 1973.
- Graham *et al.*, *J. Gen. Virol.*, 36:59-72, 1977.
- 5 Gregoriadis (ed.), In: *Drug Carriers in Biology and Medicine*, pp 287-341, 1979.
- Grunhaus and Horwitz, *Seminar in Virology*, 3:237-252, 1992.
- Hancock and Lehrer, 16:82-88, 1998.
- Harland and Weintraub, *J. Cell Biol.*, 101:1094-1099, 1985.
- Hermonat and Muzycska, *Proc. Nat'l Acad. Sci. USA*, 81:6466-6470, 1984.
- 10 Hersdorffer *et al.*, *DNA Cell Biol.*, 9:713-723, 1990.
- Herz and Gerard, *Proc. Nat'l Acad. Sci. USA*, 90:2812-2816, 1993.
- Hirata, *et al.*, *Infect. Immun.* 62 (4), 1421-1426 (1994)
- Horwich, *et al.*, *J. Virol.*, 64:642-650, 1990.
- Huttner *et al.*, *Gene*, 206:85-91, 1998.
- 15 Hwang *et al.*, *Biochem Cell Biol*, 76(2-3):235-46, 1998.
- Inouye *et al.*, *Nucleic Acids Res.*, 13:3101-3109, 1985.
- Johansson *et al.*, *J. Biol. Chem.*, 273:3718-3724, 1998.
- Jones and Shenk, *Cell*, 13:181-188, 1978.
- Kaneda, *J Biol Chem*, 264(21):12126-9, 1989.
- 20 Karlsson *et al.*, *EMBO J.*, 5:2377-2385, 1986.
- Kato *et al.*, *J. Biochem. Tokyo*, 101:207-215, 1987.
- Klaassen, In: *The Pharmacological Basis of Therapeutics*, Goodman & Gilman, Eds., Pergamon Press, 8th Ed., pp. 49-61, 1990.
- Klein *et al.*, *Nature*, 327:70-73, 1987.
- 25 Kovach *et al.*, *Gene*, 166:175-176, 1995.
- Kyte J., Doolittle R.F., *J Mol Biol*, 157(1):105-32, 1982.
- Larrick *et al.*, *Antimicrob. Agents Chemother.*, 37:2534-2539, 1993.
- Larrick *et al.* *Biochem. Biophys. Res. Commun.*, 179:170-175, 1991.
- Larrick *et al.*, *Infect. Immun.*, 63:1291-1297, 1995.
- 30 Le Gal La Salle *et al.*, *Science*, 259:988-990, 1993.
- Lehrer *et al.*, *Annu. Rev. Immunol.*, 11:105-128, 1993.

- Lehrer et al., *J. Immunol. Methods*, 137:167-173.
- Levrero et al., *Gene*, 101:195-202, 1991.
- Lorian, *In: Antibiotics in Laboratory Medicine*, Satterfield (Ed), Williams & Wilkins, Philadelphia, pp. 558-559;718, 1991.
- 5 Luo and Baldwin, *Biochemistry*, 36: 8413-21, 1997.
- Macejak and Sarnow, *Nature*, 353:90-94, 1991.
- Mahoney et al., *FEBS Lett*, 377:519-522, 1995.
- Mahoney et al., *FEBS Letters*, 377, 319-322, 1995.
- Maloy and Kari, *Biopolymers*, 37:105-122, 1995.
- 10 Markowitz et al., *J. Virol.*, 62:1120-1124, 1988.
- Martin et al., *Leuko. Biol.*, 58:128-136, 1995.
- Mason et al., *Antimicrob. Agents Chemother.*, 41:624-629, 1997.
- Mulligan, *Science*, 260:926-932, 1993.
- Nicolas and Rubenstein, *In: Vectors: A survey of molecular cloning vectors and their*
- 15 *uses*, Rodriguez and Denhardt (eds.), Stoneham: Butterworth, pp. 494-513, 1988.
- Nicolau and Sene, *Biochem. Biophys. Acta*, 721:185-190, 1982.
- Nicolau et al., *Methods Enzymol.*, 149:157-176, 1987.
- Oren et al., *Biopolymers*, 47(6):451-63, 1998.
- Ostreicher, *NY State Dent. J.*, 60(3):47-49, 1994.
- 20 Paskind et al., *Virology*, 67:242-248, 1975.
- Pelletier and Sonenberg, *Nature*, 334:320-325, 1988.
- Perales et al., *Proc. Nat'l Acad. Sci.*, 91:4086-4090, 1994.
- Perez-Mendez et al., *Eur. J. Biochem.*, 256:570-579, 1998.
- Popsueva et al., *FEBS Lett.*, 391:5-8, 1996.
- 25 Porter et al., *Infect. Immun.*, 65:2396-2401, 1997.
- Potter et al., *Proc. Nat'l Acad. Sci. USA*, 81:7161-7165, 1984.
- Publications, Bern, Switzerland, pp. 555-67, 1981.
- Racher et al., *Biotechnology Techniques*, 9:169-174, 1995.
- Ragot et al., *Nature*, 361:647-650, 1993.
- 30 Reese and Betts, *In: A Practical Approach to Infectious Diseases*, (3rd ed.), Boston, Little Brown, 1991.



- Renan, *Radiother. Oncol.*, 19:197-218, 1990.
- Rich *et al.*, *Hum. Gene Ther.*, 4:461-476, 1993.
- Ridgeway, In: Rodriguez RL, Denhardt DT, ed. *Vectors: A survey of molecular cloning vectors and their uses*, Stoneham: Butterworth, pp. 467-492, 1988.
- 5 Rippe *et al.*, *Mol. Cell Biol.*, 10:689-695, 1990.
- Rosenfeld *et al.*, *Cell*, 68:143-155, 1992.
- Rosenfeld *et al.*, *Science*, 252:431-434, 1991.
- Roux *et al.*, *Proc. Nat'l Acad. Sci. USA*, 86:9079-9083, 1989.
- Russel *et al.*, *Infect Immun*, 64(5):1565-8, 1996.
- 10 Selsted *et al.*, "Purification and antibacterial activity of antimicrobial peptides of rabbit granulocytes," *Infect. Immun.*, 45:150-154, 1984.
- Sheppard *et al.*, *Journal of Chemical Society*, p.538, 1981.
- Singh *et al.*, *PNAS*, 95:14961-5, 1998.
- Stratford-Perricaudet and Perricaudet, In: *Human Gene Transfer*, Eds, O. Cohen-Haguenauer and M. Boiron, Editions John Libbey Eurotext, France, pp. 51-61, 1991.
- 15 Stratford-Perricaudet *et al.*, *Hum. Gene Ther.*, 1:241-256, 1990.
- Szoka and Papahadjopoulos, *Proc. Nat'l Acad. Sci. USA*, 75: 4194-4198, 1978.
- Temin, In: *Gene Transfer*, Kucherlapati (ed.), New York: Plenum Press, pp. 149-188, 20 1986.
- Top *et al.*, *J. Infect. Dis.*, 124:155-160, 1971.
- Tossi *et al.*, *FEBS Lett.*, 339:108-112, 1994.
- Travis *et al.*, *Am. J. Resp. Pul. Mol. Biol.* 20:872-9, 1999.
- Travis, J., *Science*, 264(5157):360-2, 1994.
- 25 Tur-Kaspa *et al.*, *Mol. Cell Biol.*, 6:716-718, 1986.
- Turner *et al.*, "Activities of LL-37, a cathelin-associated antimicrobial peptide of human neutrophils," *Antimicrob Agents Chemother*, 42:2206-2214, 1998.
- U.S. Patent 4,554,101
- Valore *et al.*, 101:1633-1642, 1998.
- 30 Varmus *et al.*, *Cell*, 25:23-36, 1981.
- Wong *et al.*, *Gene*, 10:87-94, 1980.

- Wu & Wu, *Biochemistry*, 27:887-892, 1988.
- Wu & Wu, *J. Biol. Chem.*, 262:4429-4432, 1987.
- Wu and Wu, *Adv. Drug Delivery Rev.*, 12:159-167, 1993.
- Yang *et al.*, *Proc. Nat'l Acad. Sci USA*, 87:9568-9572, 1990
- 5 Zak and Sande, *In: Action of Antibiotics in Patients*, Sabath, Ed., Hans Huber
- Zanetti *et al.*, *FEBS Lett.*, 374:1-5, 1995.
- Zanetti *et al.*, *Ann. NY Acad. Sci.* 832:147-62, 1997.
- Zelenin *et al.*, *FEBS Lett.*, 280:94-96, 1991.

**WHAT IS CLAIMED IS:**

1. An isolated antimicrobial peptide comprising the amino acid sequence:
  - 5 KNLRRIRKIIHIIKKYG-NH<sub>2</sub> (SEQ ID NO: 1),  
 KNIRRIIRKIIHIIKKYG-NH<sub>2</sub> (SEQ ID NO: 6),  
 KNIRRIIRKIIHIIKKYG (SEQ ID NO: 7),  
 KNLRRIRKIIHIIKKYG (SEQ ID NO: 8),  
 NLRRIIRKIIHIIKKY (SEQ ID NO 9),  
 10 NIRRIIRKIIHIIKKY (SEQ ID NO: 10),  
 LRRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 11),  
 LRRIIRKIIHIIKK (SEQ ID NO: 12),  
 IRRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 13),  
 IRRIIRKIIHIIKK (SEQ ID NO: 14),  
 15 LRRIIRKIIHIIK-NH<sub>2</sub> (SEQ ID NO: 15),  
 RRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 16),  
 RRIIRKIIHIIK-NH<sub>2</sub> (SEQ ID NO: 17),  
 GLRKRLRKFRNKIKEKLKKIG (SEQ ID NO: 19),  
 KRLRKFRNKIKEKLKKIG (SEQ ID NO: 20),  
 20 RKRLRKFRNKIKEKLKKIGQKI (SEQ ID NO: 21),  
 LRKFRNKIKEKLKKIGQKI (SEQ ID NO: 22),  
 LRKFRNKIKEKLKKIGQKIQG (SEQ ID NO: 23),  
 RKFRNKIKEKLKKIG (SEQ ID NO: 24),  
 KIKEKLKKIGQKIQG (SEQ ID NO: 25),  
 25 KIKEKLKKIGQKIQGLL (SEQ ID NO: 26),  
 RGLRRLGRKIAHGVKKYGPTVLRIRIA-NH<sub>2</sub> (SEQ ID NO. 27), or  
 KNLRRIRKIIHIIKKYGPTILRIIRIIG-NH<sub>2</sub> (SEQ ID NO. 28).
2. A nucleic acid molecule encoding the antimicrobial peptide comprising the amino  
 30 acid sequence:

KNLRRIIRKIIHIIKKYG-NH<sub>2</sub> (SEQ ID NO: 1),  
 KNIRRIIRKIIHIIKKYG-NH<sub>2</sub> (SEQ ID NO: 6),  
 KNIRRIIRKIIHIIKKYG- (SEQ ID NO: 7),  
 KNLRRIIRKIIHIIKKYG (SEQ ID NO: 8),  
 5 NLRRIIRKIIHIIKKY (SEQ ID NO 9),  
 NIRRIIRKIIHIIKKY (SEQ ID NO: 10),  
 LRRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 11),  
 LRRIIRKIIHIIKK (SEQ ID NO: 12),  
 IRRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 13),  
 10 IRRIIRKIIHIIKK (SEQ ID NO: 14),  
 LRRIIRKIIHIIK-NH<sub>2</sub> (SEQ ID NO: 15),  
 RRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 16),  
 RRIIRKIIHIIK-NH<sub>2</sub> (SEQ ID NO: 17)  
 GLRKRLRKFRNKIKEKLKKIG (SEQ ID NO: 19),  
 15 KRLRKFRNKIKEKLKKIG (SEQ ID NO: 20),  
 RKRLRKFRNKIKEKLKKIGQKI (SEQ ID NO: 21),  
 LRKFRNKIKEKLKKIGQKI (SEQ ID NO: 22),  
 LRKFRNKIKEKLKKIGQKIQG (SEQ ID NO: 23),  
 RKFRNKIKEKLKKIG (SEQ ID NO: 24),  
 20 KIKEKLKKIGQKIQG (SEQ ID NO: 25),  
 KIKEKLKKIGQKIQGLL (SEQ ID NO: 26),  
 RGLRRLGRKIAHGVKKYGPTVLRIRIA-NH<sub>2</sub> (SEQ ID NO. 27), or  
 KNLRRIIRKIIHIIKKYGPTILRIIRIIG-NH<sub>2</sub> (SEQ ID NO. 28).

25 3. A pharmaceutical composition wherein said composition comprises the  
 antimicrobial peptide comprising the amino acid sequence:

KNLRRIIRKIIHIIKKYG-NH<sub>2</sub> (SEQ ID NO: 1),  
 KNIRRIIRKIIHIIKKYG-NH<sub>2</sub> (SEQ ID NO: 6),  
 30 KNIRRIIRKIIHIIKKYG (SEQ ID NO: 7),  
 KNLRRIIRKIIHIIKKYG (SEQ ID NO: 8),

5 NLRRIIRKIIHIIKKY (SEQ ID NO 9),  
 NIRRIIRKIIHIIKKY (SEQ ID NO: 10),  
 LRRIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 11),  
 LRRIRKIIHIIKK (SEQ ID NO: 12),  
 10 IRRIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 13),  
 IRRIRKIIHIIKK (SEQ ID NO: 14),  
 LRRIRKIIHIIK-NH<sub>2</sub> (SEQ ID NO: 15),  
 RRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 16),  
 RRIIRKIIHIIK-NH<sub>2</sub> (SEQ ID NO: 17),  
 15 GLRKRLRKFRNKIKEKLKKIG (SEQ ID NO: 19),  
 KRLRKFRNKIKEKLKKIG (SEQ ID NO: 20),  
 RKRLRKFRNKIKEKLKKIGQKI (SEQ ID NO: 21),  
 LRKFRNKIKEKLKKIGQKI (SEQ ID NO: 22),  
 LRKFRNKIKEKLKKIGQKIQG (SEQ ID NO: 23),  
 20 RKFRNKIKEKLKKIG (SEQ ID NO: 24),  
 KIKEKLKKIGQKIQG (SEQ ID NO: 25),  
 KIKEKLKKIGQKIQGLL (SEQ ID NO: 26),  
 RGLRRLGRKIAHGVKKYGPTVLRIRIA-NH<sub>2</sub> (SEQ ID NO. 27), or  
 KNLRRIIRKIIHIIKKYGPITLRIRIIG-NH<sub>2</sub> (SEQ ID NO. 28); and  
 25 a pharmaceutically acceptable carrier.

4. A method of inhibiting microbial growth in an environment capable of sustaining  
 such growth comprising administering to said environment a first antimicrobial peptide  
 25 comprising the amino acid sequence:

30 KNLRRIIRKIIHIIKKYG-NH<sub>2</sub> (SEQ ID NO: 1),  
 KNIRRIIRKIIHIIKKYG-NH<sub>2</sub> (SEQ ID NO: 6),  
 KNIRRIIRKIIHIIKKYG (SEQ ID NO: 7),  
 KNLRRIIRKIIHIIKKYG (SEQ ID NO: 8),  
 NLRRIIRKIIHIIKKY (SEQ ID NO 9),

- NIRRIIRKIIHIIKKY (SEQ ID NO: 10),  
 LRRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 11),  
 LRRIIRKIIHIIKK (SEQ ID NO: 12),  
 IRRRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 13),  
 5 IRRRIIRKIIHIIKK (SEQ ID NO: 14),  
 LRRIIRKIIHIIK-NH<sub>2</sub> (SEQ ID NO: 15),  
 RRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 16),  
 RRIIRKIIHIIK-NH<sub>2</sub> (SEQ ID NO: 17),  
 GLRKRLRKFRNKIKEKLKKIG (SEQ ID NO: 19),  
 10 KRLRKFRNKIKEKLKKIG (SEQ ID NO: 20),  
 RKRLRKFRNKIKEKLKKIGQKI (SEQ ID NO: 21),  
 LRKFRNKIKEKLKKIGQKI (SEQ ID NO: 22),  
 LRKFRNKIKEKLKKIGQKIQG (SEQ ID NO: 23),  
 RKFRNKIKEKLKKIG (SEQ ID NO: 24),  
 15 KIKEKLKKIGQKIQG (SEQ ID NO: 25),  
 KIKEKLKKIGQKIQGLL (SEQ ID NO: 26),  
 RGLRRLGRKIAHGVKKYGPTVLRIRIA-NH<sub>2</sub> (SEQ ID NO. 27), or  
 KNLRRIIRKIIHIIKKYGPTILRIIRIIG-NH<sub>2</sub> (SEQ ID NO. 28)..
- 20 5. The method of claim 4, wherein said first antimicrobial peptide is delivered in a pharmaceutical composition, said pharmaceutical composition, including a pharmaceutically acceptable carrier.
- 25 6. The method of claim 4, further comprising administering a second antimicrobial agent.
7. The method of claim 6, wherein said first antimicrobial peptide is administered before said second antimicrobial agent.
- 30 8. The method of claim 6, wherein said first antimicrobial peptide and said second antimicrobial agent are administered together.

9. The method of claim 6, wherein said first antimicrobial peptide is administered after said second antimicrobial agent.

5 10. The method of claim 6, wherein said second antimicrobial agent is selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO:18, SEQ ID NO: 19, SEQ ID NO: 20, SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 26, SEQ ID NO: 27 and  
10 SEQ ID NO:28, wherein the second microbial agent is a different than the first antimicrobial peptide.

11. The method of claim 6, wherein said second antimicrobial agent is selected from the group consisting of a protein synthesis inhibitor, a cell wall growth inhibitor, a cell membrane synthesis inhibitor, a nucleic acid synthesis inhibitor, and a competitive  
15 inhibitor.

12. A method of inhibiting microbial growth in a host, comprising administering to said host a first antimicrobial peptide comprising the amino acid sequence:

20 KNLRRIRKIIHIIKKYG-NH<sub>2</sub> (SEQ ID NO: 1) ,  
RGLRRLGRKIAHGVKKYGPTVLRIIRIAG (SEQ ID NO: 2),  
KIAHGVKKYGPTVLRIIRIAG (SEQ ID NO 3),  
LGRKIAHGVKKYGPTVLRII (SEQ ID NO: 4),  
RGLRRLGRKIAHGVKKYG (SEQ ID NO: 5),  
25 KNIRRIIRKIIHIIKKYG-NH<sub>2</sub> (SEQ ID NO: 6),  
KNIRRIIRKIIHIIKKYG (SEQ ID NO: 7),  
KNLRRIRKIIHIIKKYG (SEQ ID NO: 8),  
NLRRIIRKIIHIIKKY (SEQ ID NO 9),  
NIRRIIRKIIHIIKKY (SEQ ID NO: 10),  
30 LRRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 11),  
LRRIIRKIIHIIKK (SEQ ID NO: 12),



5      IRRIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 13),  
       IRRIRKIIHIIKK (SEQ ID NO: 14),  
       LRRIIRKIIHIIK-NH<sub>2</sub> (SEQ ID NO: 15),  
       RRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 16),  
       RRIIRKIIHIIK-NH<sub>2</sub> (SEQ ID NO: 17),  
       GLRKRLRKFRNKIKEKLKKIGQKIQGLLPKLAPRTDY (SEQ ID NO: 18),  
       GLRKRLRKFRNKIKEKLKKIG (SEQ ID NO: 19),  
       KRLRKFRNKIKEKLKKIG (SEQ ID NO: 20),  
       RKRLRKFRNKIKEKLKKIGQKI (SEQ ID NO: 21),  
 10     LRKFRNKIKEKLKKIGQKI (SEQ ID NO: 22),  
       LRKFRNKIKEKLKKIGQKIQG (SEQ ID NO: 23),  
       RKFRNKIKEKLKKIG (SEQ ID NO: 24),  
       KIKEKLKKIGQKIQG (SEQ ID NO: 25),  
       KIKEKLKKIGQKIQGLL (SEQ ID NO: 26),  
 15     RGLRRLGRKIAHGVKKYGPTVLRIRIA-NH<sub>2</sub> (SEQ ID NO. 27), or  
       KNLRRIIRKIIHIIKKYGPTILRIIRIIG-NH<sub>2</sub> (SEQ ID NO. 28).).

13.    A method of inhibiting the growth of drug-resistant microbial strains comprising  
 administering to an environment capable of sustaining such growth a first antimicrobial  
 20    peptide comprising the amino acid sequence:

      KNLRRIIRKIIHIIKKYG-NH<sub>2</sub> (SEQ ID NO: 1) ,  
       RGLRRLGRKIAHGVKKYGPTVLRIRIAG (SEQ ID NO: 2),  
       KIAHGVKKYGPTVLRIRIAG (SEQ ID NO 3),  
 25     LGRKIAHGVKKYGPTVLRIRI (SEQ ID NO: 4),  
       RGLRRLGRKIAHGVKKYG (SEQ ID NO: 5),  
       KNIRRIIRKIIHIIKKYG-NH<sub>2</sub> (SEQ ID NO:6),  
       KNIRRIIRKIIHIIKKYG (SEQ ID NO: 7),  
       KNLRRIIRKIIHIIKKYG (SEQ ID NO: 8),  
 30     NLRRIRKIIHIIKKY (SEQ ID NO 9),  
       NIRRIIRKIIHIIKKY (SEQ ID NO: 10),

- LRRIRKIIHIIKK-NH, (SEQ ID NO: 11),  
 LRRIRKIIHIIKK (SEQ ID NO: 12),  
 IRRIRKIIHIIKK-NH, (SEQ ID NO: 13),  
 IRRIRKIIHIIKK (SEQ ID NO: 14),  
 5 LRRIRKIIHIIK-NH, (SEQ ID NO: 15),  
 RRIIRKIIHIIKK-NH, (SEQ ID NO: 16),  
 RRIIRKIIHIIK-NH, (SEQ ID NO: 17),  
 GLRKRLRKFRNKIKEKLKKIGQKIQGLLPKLAPRTDY (SEQ ID NO: 18),  
 GLRKRLRKFRNKIKEKLKKIG (SEQ ID NO: 19),  
 10 KRLRKFRNKIKEKLKKIG (SEQ ID NO: 20),  
 RKRLRKFRNKIKEKLKKIGQKI (SEQ ID NO: 21),  
 LRKFRNKIKEKLKKIGQKI (SEQ ID NO: 22),  
 LRKFRNKIKEKLKKIGQKIQG (SEQ ID NO: 23),  
 RKFRNKIKEKLKKIG (SEQ ID NO: 24),  
 15 KIKEKLKKIGQKIQG (SEQ ID NO: 25),  
 KIKEKLKKIGQKIQGLL (SEQ ID NO: 26),  
 RGLRRLGRKIAHGVKKYGPTVLRIRIA-NH<sub>2</sub> (SEQ ID NO. 27), or  
 KNLRRIIRKIIHIIKKYGPTILRIIRIIG-NH<sub>2</sub> (SEQ ID NO. 28).
- 20 14. The method of claim 13, wherein said drug-resistant microbial strain is selected from the group consisting of *Psuedomonas aeruginosa*, *Burkholderia cepacia*, *Alcaligenes*, and *Xanthamonas*.
- 25 15. The method of claim 13, further comprising administering a second antimicrobial agent.
16. The method of claim 15, wherein said first antimicrobial peptide is administered before said second antimicrobial agent.
- 30 17. The method of claim 15, wherein said first antimicrobial peptide and said second antimicrobial agent are administered together.

18. The method of claim 15, wherein said first antimicrobial peptide is administered after said second antimicrobial agent.

5 19. The method of claim 15, wherein said second antimicrobial agent is selected from the group consisting of a protein synthesis inhibitor, a cell wall growth inhibitor, a cell membrane synthesis inhibitor, a nucleic acid synthesis inhibitor, and a competitive inhibitor.

10 20. A method of inhibiting growth of a microbial strain in a host, comprising administering to said host a first antimicrobial peptide comprising the amino acid sequence:

KNLRRIRKIIHIKKYG-NH<sub>2</sub> (SEQ ID NO: 1) ,  
 15 RGLRRLGRKIAHGVKKYGPTVLRIRIAG (SEQ ID NO: 2),  
 KIAHGVKKYGPTVLRIRIAG (SEQ ID NO 3),  
 LGRKIAHGVKKYGPTVLRIRI (SEQ ID NO: 4),  
 RGLRRLGRKIAHGVKKYG (SEQ ID NO: 5),  
 KNIRRIIRKIIHIKKYG-NH<sub>2</sub> (SEQ ID NO:6),  
 20 KNIRRIIRKIIHIKKYG (SEQ ID NO: 7),  
 KNLRRIRKIIHIKKYG (SEQ ID NO: 8),  
 NLRRIIRKIIHIKKY (SEQ ID NO 9),  
 NIRRIIRKIIHIKKY (SEQ ID NO: 10),  
 LRRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 11),  
 25 LRRIIRKIIHIKK (SEQ ID NO: 12),  
 IRRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 13),  
 IRRIIRKIIHIKK (SEQ ID NO: 14),  
 LRRIIRKIIHIK-NH<sub>2</sub> (SEQ ID NO: 15),  
 RRIIRKIIHIKK-NH<sub>2</sub> (SEQ ID NO: 16),  
 30 RRIIRKIIHIK-NH<sub>2</sub> (SEQ ID NO: 17),  
 GLRKRLRKFRNKIKEKLKKIGQKIQGLLPKLAPRTDY (SEQ ID NO: 18),


- 5 GLRKRLRKFRNKIKEKLKKIG (SEQ ID NO: 19),  
 KRLRKFRNKIKEKLKKIG (SEQ ID NO: 20),  
 RKRLRKFRNKIKEKLKKIGQKI (SEQ ID NO: 21),  
 LRKFRNKIKEKLKKIGQKI (SEQ ID NO: 22),  
 LRKFRNKIKEKLKKIGQKIQG (SEQ ID NO: 23),  
 RKFRNKIKEKLKKIG (SEQ ID NO: 24),  
 KIKEKLKKIGQKIQG (SEQ ID NO: 25),  
 KIKEKLKKIGQKIQGLL (SEQ ID NO: 26),  
 10 RGLRRLGRKIAHGVKKYGPTVLRIRIA-NH<sub>2</sub> (SEQ ID NO. 27), or  
 KNLRRIIRKIIHIIKKYGPTILRIIRIIG-NH<sub>2</sub> (SEQ ID NO. 28).
21. The method of claim 20, further comprising administering a second antimicrobial agent.
- 15 22. The method of claim 20, wherein said first antimicrobial peptide is administered before said second antimicrobial agent.
23. The method of claim 20, wherein said first antimicrobial peptide and said second antimicrobial agent are administered together.
- 20 24. The method of claim 20, wherein said antimicrobial peptide is administered after said second antimicrobial agent.
- 25 25. The method of claim 20 wherein said antimicrobial agent is selected from the group consisting of a protein synthesis inhibitor, a cell wall growth inhibitor, a cell membrane synthesis inhibitor, a nucleic acid synthesis inhibitor, and a competitive inhibitor.
- 30 26. A kit for use in inhibiting microbial growth in a host comprising first antimicrobial peptide comprising the amino acid sequence:

KNLRRIIRKIIHIIKK YG-NH<sub>2</sub> (SEQ ID NO: 1) ,  
 RGLRRLGRKIAHGVKKYGPTVLRIRIAG (SEQ ID NO: 2),  
 KIAHGVKKYGPTVLRIRIAG (SEQ ID NO 3),  
 LGRKIAHGVKKYGPTVLRII (SEQ ID NO: 4),  
 5 RGLRRLGRKIAHGVKKYG (SEQ ID NO: 5),  
 KNIRRIIRKIIHIIKKYG-NH<sub>2</sub> (SEQ ID NO:6),  
 KNIRRIIRKIIHIIKKYG (SEQ ID NO: 7),  
 KNLRRIIRKIIHIIKKYG (SEQ ID NO: 8),  
 NLRRIIRKIIHIIKKY (SEQ ID NO 9),  
 10 NIRRIIRKIIHIIKKY (SEQ ID NO: 10),  
 LRRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 11),  
 LRRIIRKIIHIIKK (SEQ ID NO: 12),  
 IRRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 13),  
 IRRIIRKIIHIIKK (SEQ ID NO: 14),  
 15 LRRIIRKIIHIK-NH<sub>2</sub> (SEQ ID NO: 15),  
 RRIIRKIIHIIKK-NH<sub>2</sub> (SEQ ID NO: 16),  
 RRIIRKIIHIK-NH<sub>2</sub> (SEQ ID NO: 17),  
 GLRKRLRKFRNKIKEKLKKIGQKIQGLLPKLAPRTDY (SEQ ID NO: 18),  
 GLRKRLRKFRNKIKEKLKKIG (SEQ ID NO: 19),  
 20 KRLRKFRNKIKEKLKKIG (SEQ ID NO: 20),  
 RKRLRKFRNKIKEKLKKIGQKI (SEQ ID NO: 21),  
 LRKFRNKIKEKLKKIGQKI (SEQ ID NO: 22),  
 LRKFRNKIKEKLKKIGQKIQG (SEQ ID NO: 23),  
 RKFRNKIKEKLKKIG (SEQ ID NO: 24),  
 25 KIKEKLKKIGQKIQG (SEQ ID NO: 25),  
 KIKEKLKKIGQKIQGLL (SEQ ID NO: 26),  
 RGLRRLGRKIAHGVKKYGPTVLRIRI-NH<sub>2</sub> (SEQ ID NO. 27), or  
 KNLRRIIRKIIHIIKKYGPTILRIIRIIG-NH<sub>2</sub> (SEQ ID NO. 28)

30 in a suitable container.


27. The kit of claim 26, further comprising a second antimicrobial agent.
28. The kit of claim 27, wherein said second antimicrobial agent is selected from the group consisting of a protein synthesis inhibitor, a cell wall growth inhibitor, a cell  
5 membrane synthesis inhibitor, a nucleic acid synthesis inhibitor, and a competitive inhibitor.
29. The method of claim 20, wherein said microbial strain is a bacteria.
- 10 30. The method of claim 20, wherein said microbial strain is a virus.
31. The method of claim 30, wherein said virus is HIV, HSV or EIAV.

Figure 1. Correlation of Luminescence and *P. aeruginosa* PAO1 Viability.






**Figure 2a. CAP18 Peptides: Concentration - Dependent<sup>1</sup>  
Killing of PAO1 in High and Low Salt<sup>1</sup>.**




<sup>1</sup> Determined by the luminescence-based assay.

**Figure 2b. SMAP29 Peptides: Concentration - Dependent<sup>1</sup> Killing of PAO1 in High and Low Salt<sup>1</sup>.**



<sup>1</sup> Determined by the luminescence-based assay.


**Figure 3A. Helical Conformation of CAP18 Peptides in Aqueous Trifluoroethanol and in the Presence of Lipopolysaccharide**



	Pi	0.1% LPS	40% TFE
CAP18	8.1	39.2	84.6
CAP18-21a	7.7	19.2	97.0
CAP18-18	3.8	13.4	64.8
CAP18-17	3.1	14.6	68.8
CAP18-15a	2.7	11.5	44.7

Helical content of CAP18 peptides. Peptides were analyzed by CD spectroscopy, and % helical content was calculated.


**Figure 3B. Helical Conformation of SMAP29 Peptides in Aqueous Trifluoroethanol and in the Presence of Lipopolysaccharide**



	Pi	0.1% LPS	40% TFE
SMAP29	12.6	30.1	80.9
SMAP29-18	2.0	16.4	45.4
SMAP29X-18A	45.6	-	88.8
SMAP29X-14A	13.9	-	77.4

Helical content of SMAP29 peptides. Peptides were analyzed by CD spectroscopy, and % helical content was calculated.

**Figure 4. A Hydrophobicity Gradient Correlates with Antibacterial Potency of the  $\alpha$ -Helical Cathelicidin Peptides.**



**Figure 5a. Effect of CAP18 peptides on Human Red Blood Cells**




Figure 5b Effect of SMAP29 Peptides







Figure 6. SMAP29 is Efficacious in an *In Vivo* Ovine Model of Acute Pneumonia



## Major Conformer A of Smap29 by 2D NMR



FIG. 7

## Major Conformer B of Smap29 by 2D NMR




FIG. 8

Antimicrobial Activity of SMAP-28 in 100 mM NaCl  
(Gram-Positive Bacteria)




FIG. 9

**Antimicrobial Activity of SMAP-28 in 100 mM NaCl  
(Gram-Negative Bacteria)**




FIG. 10

## SEQUENCE LISTING

<110> TACK, BRIAN  
MCCRAY, PAUL  
WELSH, MICHAEL  
TRAVIS, SUE M.  
LEHRER, ROBERT

<120> ALPHA HELICAL PEPTIDES WITH BROAD SPECTRUM  
ANTIMICROBIAL ACTIVITY THAT ARE INSENSITIVE TO SALT

<130> IOWA:026WO

<140> UNKNOWN

<141> 2000-08-18

<150> 60/149,886

<151> 1999-08-18

<160> 28

<170> PatentIn Ver. 2.0

<210> 1

<211> 18

<212> PRT

<213> Ovis aries

<400> 1

Lys Asn Leu Arg Arg Ile Ile Arg Lys Ile Ile His Ile Ile Lys Lys  
1 5 10 15

Tyr Gly

<210> 2

<211> 29

<212> PRT

<213> Ovis aries

<400> 2

Arg Gly Leu Arg Arg Leu Gly Arg Lys Ile Ala His Gly Val Lys Lys  
1 5 10 15

Tyr Gly Pro Thr Val Leu Arg Ile Ile Arg Ile Ala Gly  
20 25

<210> 3

<211> 21

<212> PRT

<213> Ovis aries

&lt;400&gt; 3

Lys Ile Ala His Gly Val Lys Lys Leu Gly Pro Thr Val Leu Arg Ile  
1 5 10 15

Ile Arg Ile Ala Gly  
20

&lt;210&gt; 4

&lt;211&gt; 20

&lt;212&gt; PRT

&lt;213&gt; Ovis aries

&lt;400&gt; 4

Leu Gly Arg Lys Ile Ala His Gly Val Lys Lys Tyr Gly Pro Thr Val  
1 5 10 15

Leu Arg Ile Ile  
20

&lt;210&gt; 5

&lt;211&gt; 18

&lt;212&gt; PRT

&lt;213&gt; Ovis aries

&lt;400&gt; 5

Arg Gly Leu Arg Arg Leu Gly Arg Lys Ile Ala His Gly Val Lys Lys  
1 5 10 15

Tyr Gly

&lt;210&gt; 6

&lt;211&gt; 20

&lt;212&gt; PRT

&lt;213&gt; Ovis aries

&lt;400&gt; 6

Lys Asn Ile Arg Arg Ile Ile Arg Lys Ile Ile His Ile Ile Lys Lys  
1 5 10 15

Tyr Gly Asn His  
20

&lt;210&gt; 7

&lt;211&gt; 18

&lt;212&gt; PRT

&lt;213&gt; Ovis aries



&lt;400&gt; 7

Lys Asn Ile Arg Arg Ile Ile Arg Lys Ile Ile His Ile Ile Lys Lys  
1 5 10 15

Tyr Gly

&lt;210&gt; 8

&lt;211&gt; 18

&lt;212&gt; PRT

&lt;213&gt; Ovis aries

&lt;400&gt; 8

Lys Asn Leu Arg Arg Ile Ile Arg Lys Ile Ile His Ile Ile Lys Lys  
1 5 10 15

Tyr Gly

&lt;210&gt; 9

&lt;211&gt; 16

&lt;212&gt; PRT

&lt;213&gt; Ovis aries

&lt;400&gt; 9

Asn Leu Arg Arg Ile Ile Arg Lys Ile Ile His Ile Ile Lys Lys Tyr  
1 5 10 15

&lt;210&gt; 10

&lt;211&gt; 16

&lt;212&gt; PRT

&lt;213&gt; Ovis aries

&lt;400&gt; 10

Asn Ile Arg Arg Ile Ile Arg Lys Ile Ile His Ile Ile Lys Lys Tyr  
1 5 10 15

&lt;210&gt; 11

&lt;211&gt; 16

&lt;212&gt; PRT

&lt;213&gt; Ovis aries

&lt;400&gt; 11

Leu Arg Arg Ile Ile Arg Lys Ile Ile His Ile Ile Lys Lys  
1 5 10 15

&lt;210&gt; 12

&lt;211&gt; 14

&lt;212&gt; PRT

&lt;213&gt; Ovis aries

&lt;400&gt; 12

Leu Arg Arg Ile Ile Arg Lys Ile Ile His Ile Ile Lys Lys  
1 5 10

&lt;210&gt; 13

&lt;211&gt; 16

&lt;212&gt; PRT

&lt;213&gt; Ovis aries

&lt;400&gt; 13

Ile Arg Arg Ile Ile Arg Lys Ile Ile His Ile Ile Lys Lys  
1 5 10 15

&lt;210&gt; 14

&lt;211&gt; 14

&lt;212&gt; PRT

&lt;213&gt; Ovis aries

&lt;400&gt; 14

Ile Arg Arg Ile Ile Arg Lys Ile Ile His Ile Ile Lys Lys  
1 5 10

&lt;210&gt; 15

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Ovis aries

&lt;400&gt; 15

Leu Arg Arg Ile Ile Arg Lys Ile Ile His Ile Ile Lys  
1 5 10 15

&lt;210&gt; 16

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Ovis aries

&lt;400&gt; 16

Arg Arg Ile Ile Arg Lys Ile Ile His Ile Ile Lys Lys  
1 5 10 15

&lt;210&gt; 17

&lt;211&gt; 14

&lt;212&gt; PRT

&lt;213&gt; Ovis aries

&lt;400&gt; 17

Arg Arg Ile Ile Arg Lys Ile Ile His Ile Ile Lys  
1 5 10

&lt;210&gt; 18

&lt;211&gt; 37

&lt;212&gt; PRT

&lt;213&gt; Oryctolagus cuniculus

&lt;400&gt; 18

Gly Leu Arg Lys Arg Leu Arg Lys Phe Arg Asn Lys Ile Lys Glu Lys  
1 5 10 15

Leu Lys Lys Ile Gly Gln Lys Ile Gln Gly Leu Leu Pro Lys Leu Ala  
20 25 30

Pro Arg Thr Asp Tyr  
35

&lt;210&gt; 19

&lt;211&gt; 21

&lt;212&gt; PRT

&lt;213&gt; Oryctolagus cuniculus

&lt;400&gt; 19

Gly Leu Arg Lys Arg Leu Arg Lys Phe Arg Asn Lys Ile Lys Glu Lys  
1 5 10 15

Leu Lys Lys Ile Gly  
20

&lt;210&gt; 20

&lt;211&gt; 18

&lt;212&gt; PRT

&lt;213&gt; Oryctolagus cuniculus

&lt;400&gt; 20

Lys Arg Leu Arg Lys Phe Arg Asn Lys Ile Lys Glu Lys Leu Lys Lys  
1 5 10 15

Ile Gly

&lt;210&gt; 21

&lt;211&gt; 22

&lt;212&gt; PRT

&lt;213&gt; Oryctolagus cuniculus

&lt;400&gt; 21

Arg Lys Arg Leu Arg Lys Phe Arg Asn Lys Ile Lys Glu Lys Leu Lys  
1 5 10 15

Lys Ile Gly Gln Lys Ile  
20

&lt;210&gt; 22

&lt;211&gt; 19

&lt;212&gt; PRT

&lt;213&gt; Oryctolagus cuniculus

&lt;400&gt; 22

Leu Arg Lys Phe Arg Asn Lys Ile Lys Glu Lys Leu Lys Lys Ile Gly  
1 5 10 15

Gln Lys Ile

<210> 23  
<211> 21  
<212> PRT  
<213> *Oryctolagus cuniculus*

<400> 23  
Leu Arg Lys Phe Arg Asn Lys Ile Lys Glu Lys Leu Lys Lys Ile Gly  
1 5 10 15

Gln Lys Ile Gln Gly  
20

<210> 24  
<211> 15  
<212> PRT  
<213> *Oryctolagus cuniculus*

<400> 24  
Arg Lys Phe Arg Asn Lys Ile Lys Glu Lys Leu Lys Lys Ile Gly  
1 5 10 15

<210> 25  
<211> 15  
<212> PRT  
<213> *Oryctolagus cuniculus*

<400> 25  
Lys Ile Lys Glu Lys Leu Lys Lys Ile Gly Gln Lys Ile Gln Gly  
1 5 10 15

<210> 26  
<211> 17  
<212> PRT  
<213> *Oryctolagus cuniculus*

<400> 26  
Lys Ile Lys Glu Lys Leu Lys Lys Ile Gly Gln Lys Ile Gln Gly Leu  
1 5 10 15

Leu

<210> 27  
<211> 30  
<212> PRT  
<213> *Ovis aries*

<400> 27  
Arg Gly Leu Arg Arg Leu Gly Arg Lys Ile Ala His Gly Val Lys Lys  
1 5 10 15

Tyr Gly Pro Thr Val Leu Arg Ile Ile Arg Ile Ala  
                   20                  25                  30

<210> 28  
 <211> 31  
 <212> PRT  
 <213> Ovis aries

<400> 28  
 Lys Asn Leu Arg Arg Ile Ile Arg Lys Ile Ile His Ile Ile Lys Lys  
   1                  5                  10                  15

Tyr Gly Pro Thr Ile Leu Arg Ile Ile Arg Ile Ile Gly  
                   20                  25                  30

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 00/22781

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K14/47 C07K7/08 A61K38/17 A61P31/04

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

WPI Data, PAJ, CHEM ABS Data, EPO-Internal, BIOSIS, MEDLINE

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	TRAVIS, SUE M. ET AL.: "BACTERICIDAL ACTIVITY OF MAMMALIAN CATHELICIDIN-DERIVED PEPTIDES" INFECT IMMUN (2000) 68(5) 2748-2755, XP002157165 the whole document	1-31
X	TOSSI ALESSANDRO ET AL: "Design of synthetic antimicrobial peptides based on sequence analogy and amphipathicity." EUROPEAN JOURNAL OF BIOCHEMISTRY, vol. 250, no. 2, December 1997 (1997-12), pages 549-558, XP000978718 ISSN: 0014-2956	1-5, 12-14, 20,29
Y	figures 1-3	6-11, 15-19, 21-25

☒ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

12 January 2001

Date of mailing of the international search report

29/01/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Cervigni, S

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/22781

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MERLUZZI, LAURA ET AL.: "BIOLOGICAL ACTIVITY OF A NOVEL CATHELICIDIN-DERIVED ANTIMICROBIAL PEPTIDE FROM SHEEP" PEPT 1996 PROC EUR PEPT SYMP 24TH (1998) MEETING DATE 1996 639-640 EDITOR(S) RAMAGE ROBERT EPTON ROGER PUBLISHER MAYFLOWER SCIENTIFIC KINGSWINFORD UK, XP002157166	1-5, 12-14, 20,29
Y	the whole document	6-11, 15-19, 21-25
X	TOSSI ET AL: "Identification and characterization of a primary antibacterial domain in CAP18, a lipopolysaccharide binding protein from rabbit leukocytes" FEBS LETTERS,NL,ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, vol. 339, no. 1/02, 1994, pages 108-112, XP002083170 ISSN: 0014-5793	1-5, 12-14, 20,29
Y	abstract; table 1	6-11, 15-19, 21-25
Y	BALS ROBERT ET AL: "The peptide antibiotic LL-37/hCAP-18 is expressed in eptelia of the human lung where it has broad antimicrobial activity at the airway surface." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES, vol. 95, no. 16, 4 August 1998 (1998-08-04), pages 9541-9546, XP002157167 Aug. 4, 1998 ISSN: 0027-8424 abstract page 9544, column 2 -page 9545, column 2	6-11, 15-19, 21-25
A	HUTTNER K M ET AL: "Localization and genomic organization of sheep antimicrobial peptide genes" GENE: AN INTERNATIONAL JOURNAL ON GENES AND GENOMES,GB,ELSEVIER SCIENCE PUBLISHERS, BARKING, vol. 206, no. 1, 5 January 1998 (1998-01-05), pages 85-91, XP004102876 ISSN: 0378-1119	



FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-31 (all partially)

Isolated antimicrobial peptides having structural features related to SMAP 29 (SEQ ID 2) and fragments thereof, pharmaceutical compositions containing them and methods of inhibiting microbial growth.

2. Claims: 1-31 (all partially)

Isolated antimicrobial peptides having structural features related to RCAP 18 (SEQ ID 18) and fragments thereof, pharmaceutical compositions containing them and methods of inhibiting microbial growth.

3. Claims: 1-31 (all partially)

Isolated synthetic antimicrobial peptides related to SMAP 29 or RCAP 18 but having alternate structural properties (SEQ IDs 1,6-17), pharmaceutical compositions containing them and methods of inhibiting microbial growth.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☒ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**